Atmospheric NLTE-Models for the Spectroscopic Analysis of Blue Stars with Winds

III. X-ray emission from wind-embedded shocks*

Luiz P. Carneiro¹, J. Puls¹, J. O. Sundqvist^{2,3}, and T. L. Hoffmann¹

¹ LMU Munich, Universitätssternwarte, Scheinerstr. 1, 81679 München, Germany, e-mail: luiz@usm.uni-muenchen.de

² Centro de Astrobiología, CSIC-INTA, Ctra. Torrejón a Ajalvir km.4, 28850 Madrid, Spain

³ Instituut voor Sterrenkunde, KU Leuven, Celestijnenlaan 200D, 3001 Leuven, Belgium

Received; Accepted

ABSTRACT

Context. X-rays/EUV radiation emitted from wind-embedded shocks in hot, massive stars can affect the ionization balance in their outer atmospheres, and can be the mechanism responsible for the production of highly ionized atomic species detected in stellar wind UV spectra.

Aims. To allow for these processes in the context of spectral analysis, we have implemented the emission from wind-embedded shocks and related physics into our unified, NLTE model atmosphere/spectrum synthesis code FASTWIND.

Methods. The shock structure and corresponding emission is calculated as a function of user-supplied parameters (volume filling factor, radial stratification of shock strength, and radial onset of emission). We account for a temperature and density stratification inside the post-shock cooling zones, calculated for radiative and adiabatic cooling in the inner and outer wind, respectively. The high-energy absorption of the cool wind is considered by adding important K-shell opacities, and corresponding Auger ionization rates have been included into the NLTE network. To test our implementation and to check the resulting effects, we calculated a comprehensive model grid with a variety of X-ray emission parameters.

Results. We tested and verified our implementation carefully against corresponding results from various alternative model atmosphere codes, and studied the effects from shock emission for important ions from He, C, N, O, Si, and P. Surprisingly, dielectronic recombination turned out to play an essential role for the ionization balance of O iv/O v in stars (particularly dwarfs) with $T_{\text{eff}} \sim 45,000$ K. Finally, we investigated the frequency dependence and *radial* behavior of the mass absorption coefficient, $\kappa_v(r)$, important in the context of X-ray line formation in massive star winds.

Conclusions. In almost all considered cases, *direct* ionization is of major influence (because of the enhanced EUV radiation field), and Auger ionization significantly affects only N vI and O vI. The approximation of a radially constant κ_{ν} is justified for $r \gtrsim 1.2R_*$ and $\lambda \lesssim 18$ Å, and also for many models at longer wavelengths. To estimate the actual *value* of this quantity, however, the He II opacities need to be calculated from detailed NLTE modeling, at least for wavelengths longer than 18 to 20 Å, and information on the individual CNO abundances has to be present.

Key words. Methods: numerical - stars: atmospheres - stars: early-type - stars: winds, outflows - X-rays: stars

1. Introduction

Most of our knowledge about the physical parameters of hot stars has been inferred by means of quantitative spectroscopy, i.e., the analysis of stellar spectra based on atmospheric models. The computation of such models is quite challenging, mostly because of the intense radiation fields of hot stars leading to various effects that are absent in the atmospheres of cooler ones, such as the requirement for a kinetic equilibrium description (also simply called NLTE = non-LTE) and the presence of strong, radiation-driven winds.

In recent decades, a number of numerical codes have been developed which enable the calculation of synthetic profiles/spectral energy distributions (SEDs) from such hot stars. Apart from plane-parallel, hydrostatic codes that can be used to analyze those atmospheres which are less affected by the wind (e.g., TLUSTY, Hubeny 1998; DETAIL/SURFACE, Giddings 1981, Butler & Giddings 1985), all of these codes apply the concept of *unified* (or *global*) model atmospheres (Gabler et al. 1989) which aims at a consistent treatment of both photosphere and wind, i.e., including (steady-state) mass loss and velocity fields. Examples of such codes are CMFGEN (Hillier & Miller 1998), PHOENIX (Hauschildt 1992), PoWR (Gräfener et al. 2002), WM-BASIC (Pauldrach et al. 2001) and FASTWIND (Puls et al. 2005, Rivero González et al. 2012a).¹ A brief comparison of these different codes can be found in Puls (2009).

In the present paper, we report on recent progress to improve the capabilities of FASTWIND, which is widely used to analyze the *optical* spectra of hot massive stars (e.g., in the context of the VLT-FLAMES survey of massive stars, Evans et al. 2008, and the VLT-FLAMES Tarantula Survey, Evans et al. 2011). One of the most challenging aspects of these surveys was the analysis of the atmospheric nitrogen content (processed in the stellar core by the CNO-cycle and transported to the outer layers by rotational mix-

^{*} Appendices A, B, and C are only available in electronic form at http://www.edpsciences.org

¹ The multi-component code developed by Krtička & Kubát (2001) that will be referred to later on has been designed to calculate the wind properties, and has not been used for diagnostic purposes so far.

ing), in order to derive stringent constraints for up-to-date evolutionary calculations. Though the optical nitrogen analysis of B-stars (dwarfs and supergiants with not too dense winds) could still be performed by a hydrostatic code (in this case TLUSTY, e.g., Hunter et al. 2007, 2008), a similar analysis of hotter stars with denser winds required the application of unified model atmospheres, due to the wind impact onto the strategic nitrogen lines (Rivero González et al. 2011, 2012a, Martins et al. 2012). Moreover, because of the complexity of the involved processes, the precision of the derived nitrogen abundances² is still questionable. To independently check this precision and to obtain further constraints, a parallel investigation of the carbon (and oxygen) abundances is urgently needed, since at least the N/C abundance ratio as a function of N/O might be predicted almost independent from the specific evolutionary scenario (Przybilla et al. 2010), and thus allows individually derived spectroscopic abundances to be tested (see also Martins et al. 2015a).

As shown by Martins & Hillier (2012), however, the optical diagnostics of carbon in O-stars is even more complex than the nitrogen analysis, since specific, important levels are pumped by a variety of UV resonance lines. Thus, an adequate treatment of UV lines (both for the optical diagnostics, but also to constrain the results by an additional analysis of carbon lines located in the UV) is inevitable. If at least part of these lines are formed in the wind, the inclusion of X-ray and EUV emission from windembedded shocks turns out to be essential (see below); this is the main reason (though not the only one) for our current update of FASTWIND. Other codes such as CMFGEN, PoWR, and WM-BASIC already include these processes, thus enabling the modeling of the UV (e.g., Pauldrach et al. 2001, Crowther et al. 2002, Hamann & Oskinova 2012) and the analysis of carbon (plus nitrogen and oxygen, e.g., Bouret et al. 2012, Martins et al. 2015a,b for the case of Galactic O-stars).

X-ray emission from hot stars has been measured at soft (0.1 to \geq 2 keV) and harder energies, either at low resolution in the form of a quasi-continuum, or at high resolution allowing the investigation of individual lines (e.g., Oskinova et al. 2006, Owocki & Cohen 2006, Hervé et al. 2013, Leutenegger et al. 2013b, Cohen et al. 2014b, Rauw et al. 2015). Already the first X-ray satellite observatory, EINSTEIN, revealed that O-stars are soft X-ray sources (Harnden et al. 1979, Seward et al. 1979), and Cassinelli & Swank (1983) were the first to show that the observed X-ray emission is due to thermal emission, dominated by lines. Follow-up investigations, particularly by ROSAT, have subsequently allowed us to quantify X-ray properties for many OB-stars (see Kudritzki & Puls 2000 and references therein). Accounting also for more recent work based on CHANDRA and XMM-NEWTON, it was found that the intrinsic X-ray emission of 'normal' O-stars is highly constant w.r.t. time (e.g., Nazé et al. 2013), and that the level of X-ray emission is quite strictly related to basic stellar and wind parameters, e.g., $L_x/L_{bol} \approx 10^{-7}$ for O-stars (Chlebowski et al. 1989, Sana et al. 2006, Nazé et al. 2011).

Such X-ray emission is widely believed to originate from wind-embedded shocks, and to be related to the line-driven instability (LDI, e.g., Lucy & Solomon 1970, Owocki & Rybicki 1984, Owocki et al. 1988, Owocki 1994, Feldmeier 1995). In terms of a stationary description, a simple model (e.g., Hillier et al. 1993, Cassinelli et al. 1994) assumes randomly distributed shocks above a minimum radius, $R_{\min} \approx 1.5 R_*$ (consistent with X-ray line diagnostics, e.g., Leutenegger et al. 2013b, but see also Rauw et al. 2015) where the hot shocked gas (with temperatures of a few million Kelvin and a volume filling factor on the order of 10^{-3} to a few 10^{-2}) is collisionally ionized/excited and emits X-ray/EUV photons due to spontaneous decay, radiative recombinations and bremsstrahlung. The ambient, *cool* wind then re-absorbs part of the emission, mostly via K-shell processes. The strength of this wind-absorption has a strong frequency dependence. For energies beyond 0.5 keV (e.g., the CHANDRA-bandpass), the absorption is quite modest (e.g., Cohen et al. 2011), whilst for softer X-rays and the EUV regime the absorption is significant, even for winds with low mass-loss rate (e.g., Cohen et al. 1996). In the latter case, only a small fraction of the produced radiation actually leaves the wind.

This simple model, sometimes extended to account for the post-shock cooling zones of radiative and adiabatic shocks (see Feldmeier et al. 1997a, but also Owocki et al. 2013), is used in the previously mentioned NLTE codes, particularly to account for the influence of X-ray/EUV emission on the photo-ionization rates.

Since the detection of high ionization stages in stellar wind UV spectra, such as O vi, S vi, and N v (Snow & Morton 1976, Lamers & Morton 1976, Lamers & Rogerson 1978), that cannot be produced in a cool wind (thus, denoted by 'superionization'), the responsible mechanism was (and partly still is) subject to debate. Because the X-ray and associated EUV luminosity emitted by the shocks is quite strong, it can severely affect the degree of ionization of highly ionized species, by Auger ionization (Macfarlane et al. 1993), and even more by direct ionization in the EUV (Pauldrach et al. 1994, 2001). A first *systematic* investigation of stellar parameters, mass loss, and X-ray luminosity has been performed by Garcia (2005).

In this paper, we present our approach for implementing wind-embedded shocks into FASTWIND, to allow for further progress as outlined above, and report on corresponding tests and first results. In Sect. 2, our model for the X-ray emission and cool-wind absorption is described, together with the coupling to the equations of statistical equilibrium. In Sect. 3 we present our model grid which constitutes the basis of our further discussion. Sect. 4 provides some basic tests, and Sect. 5 presents first results. In particular, we discuss how the ionization fractions of specific, important ions are affected by X-ray emission, and how these fractions change when the description of the emission (filling factors, shock temperatures) is varied (Sect. 5.1). We compare with results from other studies (Sect. 5.1.4), and investigate the impact of Auger compared to direct ionization (Sect. 5.2). We discuss the impact of dielectronic recombination in Ov in Sect. 5.3, and comment on the radial behavior of the mass absorption coefficient (as a function of wavelength), an important issue for X-ray line diagnostics (Sect. 5.4). Finally, we present our summary and conclusions in Sect. 6.

2. Implementation of X-ray emission and absorption in FASTWIND

Our implementation of the X-ray emission and absorption from wind-embedded shocks follows closely the implementation by Pauldrach et al. (2001) (for WM-BASIC, see also Pauldrach et al. 1994), which in turn is based on the model for shock cooling zones developed by Feldmeier et al. (1997a) (see Sect. 1). Except for the description of the cooling zones, this implementation is similar to the approaches by Hillier & Miller (1998) (CMFGEN) (who use a different definition of the filling factor, see below),

² which, for early-type O-stars, suggest very efficient mixing processes already at quite early stages (Rivero González et al. 2012b)

Oskinova et al. (2006) (POWR), and Krtička & Kubát (2009, hereafter KK09). In the following, we summarize our approach.

2.1. X-ray Emission

Following Feldmeier et al. (1997a), the energy (per unit of volume, time and frequency), emitted by the *hot* gas into the full solid angle 4π can be written as³

$$\epsilon_{\nu} = f_{\rm X}(r)n_{\rm p}(r)n_{\rm e}(r)\Lambda_{\nu}(n_{\rm e}(r), T_{\rm s}(r)) \tag{1}$$

where $n_p(r)$ and $n_e(r)$ are the proton and electron density of the (quasi-)stationary, 'cool' (pre-shock) wind, $T_s(r)$ is the shock temperature, and $f_X(r)$ the filling factor related to the (volume) fraction of the X-ray emitting material.⁴ Indeed, this definition differs from the formulation suggested by Hillier et al. (1993, their Eq. 2), since we include here their factor⁵ 16 into f_X . This definition is then identical with that used in WM-BASIC, POWR (presumably⁶) and by KK09, whilst the relation to the filling factor used in CMFGEN, e_s , is given by

$$f_{\rm X} = 16 \, e_{\rm s}^2. \tag{2}$$

In principle, Λ_{ν} is the frequency dependent volume emission coefficient ('cooling function') per proton and electron, calculated here using the Raymond-Smith code (Raymond & Smith 1977, see also Smith et al. 2001), with abundances from the FASTWIND input, and neglecting the weak dependence on n_e . We evaluate the cooling function at a fixed electron density, $n_e = 10^{10}$ cm⁻³ (as also done, e.g., by Hillier et al. 1993 and Feldmeier et al. 1997a), and have convinced ourselves of the validity of this approximation. We note here that the only spectral features with a significant dependence on electron density are the forbidden and intercombination lines of He-like emission complexes, and even there (i) the density dependence is 'swamped' by the dependence on UV photo-excitation, and (ii) in any case the flux of the forbidden plus intercombination line complex (f+i lines are very closely spaced) is conserved.

Contrasted to the assumption of a hot plasma with a fixed post-shock temperature and density (as adopted in some of the above codes), in our implementation we account for a temperature and density stratification in the post-shock cooling zones, noting that the decreasing temperature and increasing density should significantly contribute to the shape of the emitted X-ray spectrum (Krolik & Raymond 1985). To this end, we adopt the structure provided by Feldmeier et al. (1997a), and integrate the emitted energy (Eq. 1) over the cooling zone,

$$\overline{\epsilon}_{\nu} = f_{\rm X}(r) n_{\rm p}(r) n_{\rm e}(r) \overline{\Lambda_{\nu}} (10^{10} \,{\rm cm}^{-3}, T_{\rm s}(r)), \tag{3}$$

with

$$\overline{\Lambda_{\nu}}(T_{\mathrm{s}}(r)) = \pm \frac{1}{L_c} \int_r^{r \pm L_c} f^2(r') \Lambda_{\nu}(T_{\mathrm{s}}(r) \cdot g(r')) \,\mathrm{d}r', \quad (4)$$

where *r* is the position of the shock front, and L_c the spatial extent of the cooling zone. In this formulation, the '+' sign corresponds to a reverse shock, and the '-' sign to a forward one. The functions *f* and *g* provide the normalized density and temperature stratification inside the cooling zone, and are calculated

following Feldmeier et al. (1997a), accounting for radiative and adiabatic cooling in the inner and outer wind, respectively (see Sect. 2.3). We integrate over 1,000 subgrid points within L_c , finding identical results for both f(r) and g(r) as well as for $\overline{\Lambda_v}$, compared to the original work (Figs. 1/7 and 2/8 in Feldmeier et al. 1997a). Note that by setting f = g = 1, we are able to return to non-stratified, isothermal shocks.

In our implementation, the (integrated) cooling function and thus the emissivity is evaluated in the interval between 1 eV and 2.5 keV, for a bin-size of 2.5 eV. These emissivities are then resampled onto our coarser frequency grid as used in FASTWIND, in such a way as to preserve $\int \varepsilon_{\nu} d\nu$ in each of the coarser subintervals, thus enabling correct photo-integrals for the rate equations.

The immediate post-shock temperature, $T_s(r)$, entering Eq. 4, follows from the Rankine-Hugoniot equations:

$$T_{\rm s}(r) = \frac{3}{16} \frac{\mu m_{\rm H}}{k_{\rm B}} \left(u^2 + \left[\frac{14}{5} a_{\rm s}^2 \left(1 - \frac{3}{14} \frac{a_{\rm s}^2}{u^2} \right) \right] \right)$$
(5)

where *u* is the jump velocity, μ the mean atomic weight, and a_s the adiabatic upstream sound speed. For simplicity, we calculate the shock temperature from a more approximate expression, neglecting the term in the square bracket, i.e., assuming the strong shock scenario ($u^2 \gg a_s^2$):

$$T_{\rm s}(r) = \frac{3}{16} \frac{\mu m_{\rm H}}{k_{\rm B}} u^2 \tag{6}$$

To derive T_s , we thus need to specify the jump velocity u, adopted in accordance with Pauldrach et al. (1994, their Eq. 3) as

$$u(r) = u_{\infty} \left[\frac{v(r)}{v_{\infty}} \right]^{\gamma_x}$$
(7)

where u_{∞} is the maximum jump speed which in our implementation is an input parameter (on the order of 300 to 600 km s^{-1} , corresponding to a maximum shock temperature, $T_{\rm s}^{\infty} \approx 10^6$ to $5 \cdot 10^6$ K for O-stars), together with the exponent γ_x (in the typical range 0.5...2) that couples the jump velocity with the outflow velocity, controlling the shock strength. A parameterization such as Eq. 7 is motivated primarily by the observed socalled 'black troughs' in UV P-Cygni profiles. Namely, when modeled using a steady-state wind⁷, such black troughs can only be reproduced when assuming a velocity dispersion that increases in parallel with the outflow velocity, interpreted as a typical signature of wind-structure (e.g., Groenewegen & Lamers 1989, Haser 1995). Note, however, that Eq. 7 only represents one possible implementation of the radial distribution of windshock strengths, and that ultimately the user will be responsible for her/his choice of parameterization (see also discussion in Sect. 6).

The last required parameter is the onset radius of the Xray emission, R_{\min} . This value is controlled by two input parameters, R_{\min}^{input} and a factor m_x (the latter in accordance with Pauldrach et al. 1994). From these values, R_{\min} is calculated via

$$R_{\min} = \min(R_{\min}^{\text{input}}, r(v_{\min})) \quad \text{with} \quad v_{\min} = m_{x} a_{s} \quad (8)$$

For all radii $r > R_{min}$, the X-ray emission is switched on. R_{min} values from 1.1 to 1.5 R_* are, e.g., supported by Pauldrach et al. (1994), from their analysis of the O vi resonance

³ The corresponding emissivity is lower by a factor $1/4\pi$.

⁴ The actual, local pre-shock density may be different from its quasistationary equivalent, but this difference gets absorbed in the f_X -factor.

⁵ accounting for the density jump in a strong adiabatic shock

⁶ We were not able to find a definite statement, but Oskinova et al. (2006) also refer to Feldmeier et al. (1997a).

⁷ but see Lucy 1982, Puls et al. 1993, Sundqvist et al. 2012b for the case of time-dependent, non-monotonic velocity fields

lines. Hillier et al. (1993) analyzed the sensitivity to $R_{\rm min}$, pointing to indistinguishable X-ray-flux differences when the onset is varied between 1.5 and 2 R_* . Recent analyses of X-ray *line* emission from hot star winds also point to values around 1.5 R_* (e.g., Leutenegger et al. 2006, Oskinova et al. 2006, Hervé et al. 2013, Cohen et al. 2014b), though Rauw et al. (2015) derived a value of 1.2 R_* for the wind of λ Cep.

2.2. X-ray absorption and Auger ionization

Besides the X-ray emission, the absorption by the 'cold' background wind⁸ and needs to be computed.

In FASTWIND, the 'cool' wind opacity is computed in NLTE, and to include X-ray absorption requires that we (i) extend the frequency grid and coupled quantities (standard⁹ opacities and emissivities, radiative transfer) into the X-ray domain (until 2.5 keV \approx 5 Å), and (ii) compute the additional absorption by inner shell electrons, leading, e.g., to Auger ionization. So far, we included only K-shell absorption for light elements using data from Daltabuit & Cox (1972). L- and M-shell processes for heavy elements – which are also present in the considered energy range – have not been incorporated until now, but would lead to only marginal effects, as test calculations by means of WM-BASIC have shown.

We checked that the K-shell opacities by Daltabuit & Cox (1972) are quite similar (with typical differences less than 5%) to the alternative and more 'modern' dataset from Verner & Yakovlev (1995), at least in the considered energy range (actually, even until 3.1 keV).¹⁰

The reader may note that though the provided dataset includes K-shell opacities from the elements C, N, O, Ne, Mg, Si, and S, the last one (S) has threshold energies beyond our maximum energy, 2.5 keV, so that K-shell absorption and Auger ionization for this element is not considered in our model.

After calculating the radiative transfer in the X-ray regime, accounting for standard and K-shell opacities as well as standard and X-ray emissivities, we are able to calculate the corresponding photo-rates required to consider Auger-ionization in our NLTE treatment. Here, we do not only include the transition between ions separated by a charge-difference of two (such as, e.g., the ionization from O_{IV} to O_{VI}), but we follow Kaastra & Mewe (1993) who stressed the importance of cascade ionization processes, enabling a sometimes quite extended range of final ionization stages. E.g., the branching ratio for O IV to O V vs. O IV to O VI is quoted as 96:9904 whilst the branching ratios for Sim to Sin/Siv/Sivi are 3:775:9222, i.e., here the major Auger-ionization occurs for the process III to VI. In our implementation of Auger ionization, we have accounted for all possible branching ratios following the data provided by Kaastra & Mewe.

Finally, we re-iterate that in addition to such inner shell absorption/Auger ionization processes, direct ionization due to Xrays/enhanced EUV radiation (e.g., of Ov and Ovi) is essential and 'automatically' included in our FASTWIND modeling. The impact of direct vs. Auger ionization will be compared in Sect. 5.2.

2.3. Radiative and Adiabatic cooling

As pointed out in Sect. 2.1, the shock cooling zones are considered to be dominated by either radiative or adiabatic cooling, depending on the location of the shock front. More specifically, the transition between the two cooling regimes is obtained from the ratio between the radiative cooling time, t_c , i.e., the time required by the shocked matter to return to the ambient wind temperature, and the flow time, t_f , the time for the material to cross L_c .¹¹ In the inner part of the wind, the cooling time is shorter than the flow time, and the shocks are approximated as radiative. Further out in the wind, at low densities, $t_c \gg t_f$, and the cooling is dominated by adiabatic expansion (see also Simon & Axford 1966). In our approach, we switch from one treatment to the other when a unity ratio is reached, where $t_c/t_f \propto T_s(r)^{1.5} r v^2(r)/\dot{M}$. For typical O-supergiants and shock temperatures, the transition occurs in the outermost wind beyond $r > 50 R_*$, whilst for O-dwarfs the transition can occur at much lower radii, $r > 2.5 R_*$ or even lower for weak-winded stars.

Basically, each cooling zone is bounded by a reverse shock at the starward side and a forward shock at the outer side. Timedependent wind simulations (e.g., Feldmeier 1995) show that in the radiative case the forward shock is much weaker than the reverse one, and is thus neglected in our model. In the adiabatic case, we keep both the reverse and the forward shock, and, because of lack of better knowledge, assume equal T_s for both components ($\Theta = 1$ in the nomenclature by Feldmeier et al. 1997a), and an equal contribution of 50% to the total emission.

3. Model grid

In this section, we describe the model grid used in most of the following work. In order to allow for a grid of theoretical models that enables us to investigate different regimes of X-ray emission for different stellar types, and to perform meaningful tests, we use the same grid as presented by Pauldrach et al. (2001) (their Table 5) for discussing the predictions of their (improved) WM-BASIC code.¹² Moreover, this grid has already been used by Puls et al. (2005) to compare the results from an earlier version of FASTWIND with the WM-BASIC code.

For convenience, we present the stellar and wind parameters of this grid in Table 1. For all models, the velocity field exponent has been set to $\beta = 0.9$. Note that the FASTWIND and WM-BASIC models display a certain difference in the velocity field¹³.

All entries displayed in Table 1 refer to homogeneous winds, though for specific tests (detailed when required) we have calculated micro-clumped models as well (i.e., assuming optically thin clumps). We remind the reader that though clumping is not considered in our standard model grid, a (micro-)clumped wind could be roughly compared to our unclumped models as long as the mass-loss rate of the clumped model corresponds to the mass-loss rate of the unclumped one, divided by the square root of the clumping factor.¹⁴

⁸ The optical depths *inside* the shocked plasma are so low that absorption can be neglected there.

 $^{^9}$ = outer electron shell

¹⁰ The major reason for using data from Daltabuit & Cox (1972) was to ensure compatibility with results from WM-BASIC, to allow for meaningful comparisons. In the near future, we will update our data following Verner & Yakovlev (1995).

¹¹ Expressions for these quantities can be found in Feldmeier et al. (1997a), but see also Hillier et al. (1993).

 $^{^{12}}$ This grid, in turn is based on observational results from Puls et al. (1996), which at that time did not include the effects of wind inhomogeneities, so that the adopted mass-loss rates might be too large, by factors from ~3...6.

¹³ WM-BASIC calculates the velocity field from a consistent *hydrodynamic* approach.

¹⁴ Note, however, that K-shell opacities scale linearly with density, i.e., $\propto \dot{M}$, and as such are *not* affected by micro-clumping.

Table 1. Stellar and wind parameters of our grid models with homogeneous winds, following Pauldrach et al. (2001). For X-ray emission parameters, see text.

Model	$T_{\rm eff}$	log g	R_*	v_{∞}	М	R_{\min}				
	(kK)	$({\rm cm}{\rm s}^{-2})$	(R_{\odot})	$({\rm km}{\rm s}^{-1})$	$(10^{-6} M_{\odot} \mathrm{yr}^{-1})$	(R_*)				
Dwarfs										
D30	30	3.85	12	1800	0.008	1.24				
D35	35	3.80	11	2100	0.05	1.29				
D40	40	3.75	10	2400	0.24	1.20				
D45	45	3.90	12	3000	1.3	1.20				
D50	50	4.00	12	3200	5.6	1.23				
D55	55	4.10	15	3300	20	1.21				
Supergiants										
S30	30	3.00	27	1500	5.0	1.51				
S35	35	3.30	21	1900	8.0	1.43				
S40	40	3.60	19	2200	10	1.33				
S45	45	3.80	20	2500	15	1.25				
S50	50	3.90	20	3200	24	1.25				

All models in the present work were calculated by means of the most recent version (as described in Rivero González et al. 2012a) of the NLTE atmosphere/spectrum synthesis code FASTWIND, including the X-ray emission from windembedded shocks as outlined in Sect. 2. Let us further point out that FASTWIND calculates the temperature structure (of the photosphere and 'cold' wind) from the electron thermal balance (Kubát et al. 1999), and its major influence in the wind is via recombination rates. In most cases, this temperature structure is only slightly or moderately affected by X-ray/EUV emission, since the overall ionization balance with respect to main ionization stages¹⁵ remains rather unaffected (see Sect. 5), except for extreme X-ray emission parameters. In any case, the change of the net ionization rates for ions with edges in the soft X-ray/EUV regime is dominated by modified photo-rates (direct and Auger ionization), whilst the changes of recombination rates (due to a modified temperature) are of second order.

In FASTWIND, we used detailed model atoms for H, He, and N (described by Puls et al. 2005 and Rivero González et al. 2012a) together with C, O, P (from the WM-BASIC data base, see Pauldrach et al. 2001) and Si (see Trundle et al. 2004) as 'explicit' elements. Most of the other elements up to Zn are treated as background elements. For a description of FASTWIND and the philosophy of explicit and background elements, see Puls et al. (2005) and Rivero González et al. (2012a).

In brief, explicit elements are those used as diagnostic tools and treated with high precision, by detailed atomic models and by means of comoving frame transport for all line transitions. The background elements (i.e., the rest) are needed 'only' for the line-blocking/blanketing calculations, and are treated in a more approximate way, using parameterized ionization cross-sections following Seaton (1958) and a comoving frame transfer only for the most important lines, whilst the weaker ones are calculated by means of the Sobolev approximation.

We employed solar abundances from Asplund et al. (2009), together with a helium abundance, by number, $N_{\text{He}}/N_{\text{H}} = 0.1$.

Besides the atmospheric and wind parameters displayed in Table 1, our model of X-ray emission requires the following ad-

ditional input parameters: f_X , u_∞ , γ_x , m_x , and R_{\min}^{input} , as described in the previous section.

For most of the models discussed in Sect. 5, we calculated, per entry in Table 1, 9 different sets of X-ray emission: f_X (adopted as spatially constant) was set to 0.01, 0.03, and 0.05, whilst the maximum shock velocity, u_{∞} , was independently set to 265, 460, and 590 km s⁻¹, corresponding to maximum shock temperatures of 1, 3, and 5·10⁶ K.

For all models, we used $\gamma_x = 1.0$, $R_{\min}^{\text{input}} = 1.5 R_*$, and $m_x = 20$. This corresponds to an effective onset of X-rays, R_{\min} , between 1.2 and 1.5 R_* , or 0.1 and 0.2 v_{∞} , respectively (see Table 1, last column). Thus, our current grid comprises 9 times 11 = 99 models, and has enough resolution for comparisons with previous results from other codes and for understanding the impact of the X-ray radiation onto the ionization fractions of various elements.

4. Tests

In this section, we describe some important tests of our implementation, including a brief parameter study. A comparison to similar studies with respect to ionization fractions (also regarding the impact of Auger ionization) will be provided in Sect. 5. Of course, we have tested much more than described in the following sections, e.g.,

(i) the impact of γ_x (see also Pauldrach et al. 2001), particularly when setting γ_x to zero (and consequently forcing all shocks, independent of their position, to emit at the maximum shock temperature, T_s^{∞}). In this case and compared to our standard grid with $\gamma_x = 1$, the dwarf models cooler than 50 kK display a flux increase of 2 dex shortward of 100 Å (already for D50 this increase is barely noticeable), whilst the supergiant models display a similar increase, but for wavelengths around 10 Å and below. In terms of ionization fractions, setting γ_x to zero results in an increase of highly ionized species (e.g., O vI and N vI) by roughly one dex, from the onset of X-ray emission throughout the wind. For all other dwarf models, this increase appears only out to ~4.0 R_* . The same effect is present in the supergiant models, but for a smaller radial extent.

(ii) We compared the ionization fractions of important atoms when either treated as explicit (i.e., 'exact') or as background (i.e, approximate) elements (cf. Sect. 3), and we mostly found an excellent agreement¹⁶ for the complete model grid.

(iii) During our study on the variations of the mass absorption coefficient with $T_{\rm eff}$ and r in the X-ray regime (see Sect. 5.4), we also compared our opacities with those predicted by KK09 (their Fig. 15, displaying mass absorption coefficient vs. wavelength), and we were able to closely reproduce their results, at least shortward of 21 Å (including the dominating O rv/O v K-shell edge), but our model produces lower opacities on the longward side, thus indicating a different He ionization balance (see Sect. 5.4). When comparing the *averaged* (between 1.5 and 5 R_*) absorption coefficients in the wavelength regime shortward of 30 Å, KK09 found a slight decrease of 8% after including X-rays in their models, because of the induced ionization shift. This is consistent with our findings, which indicate, for the same range of r and λ , a decrease by 9%.

¹⁵ which dominate the heating/cooling of the cold wind plasma via corresponding free-free, bound-free and collisional (de-)excitation processes

¹⁶ In all cases, the agreement was at least satisfactory.



Fig. 1. Emergent Eddington fluxes for model S30, with $T_s^{\infty} = 3 \cdot 10^6$ K and $f_X = 0.03$, for different onset radii of X-ray emission, R_{\min} , and for a model with an unshocked wind. The vertical dotted lines refer to the He II, C II, and N III ionization edges, respectively.

4.1. Impact of various parameters

First, we study the impact of various parameters on the emergent (soft) X-ray fluxes, in particular R_{\min} , f_X , and T_s^{∞} . For these tests, we used the model S30 (see Table 1, similar to the parameters of α Cam (HD 30614, O9.5Ia)) since the latter object has been carefully investigated by Pauldrach et al. (2001, their Table 9) as well.

Before going into further details, let us clarify that the soft X-ray and EUV shock emission are composed almost entirely of narrow lines, and that the binning and blending make the spectral features look more like a pseudo-continuum, which is clearly visible in the following figures (though most of them display the emergent fluxes, and not the emissivities themselves).¹⁷

Impact of $R_{\rm min}$. The sensitivity of the X-ray fluxes on $R_{\rm min}$ is displayed in Fig. 1, where the other parameters were fixed at their center values within our small X-ray grid (i.e., $f_{\rm X} = 0.03$ and $T_{\rm s}^{\infty} = 3 \cdot 10^6 \, {\rm K}^{18}$).

Indeed, the only visible differences are present in the range between the He II edge and roughly 330 Å. Shortward of the He II edge, all fluxes are identical (though only shown down to 100 Å, to allow for a better resolution), since the (cool) wind becomes optically thick already far out in the wind at these wavelengths (He II, O IV, etc. continua, and K-shell processes). For $\lambda_{\dot{c}}$ 350 Å, on the other hand, the shock emissivity becomes too low to be of significant impact.

In this context, it is interesting to note that in ϵ CMa (B2II, the only massive hot star with EUVE data) the observed EUV emission lines in the range between 228 to 350 Å *each* have a luminosity comparable to the total X-ray luminosity in the ROSAT bandpass (Cassinelli et al. 1995), which stresses the importance of this wavelength region also from the observational side.



Fig. 2. Ratio of shock emissivity to total emissivity for model S30 from Fig. 1, with R_{\min} = 1.2 R_* . Solid line: emissivity ratio at the outer boundary, $r \approx 130 R_*$; dash-dotted: emissivity ratio at the lower boundary of X-ray emission, $r \approx 1.2 R_*$. The box located between 300 and 320 Å highlights the strong shock emissivity leading to the corresponding 'emission feature' present in Fig. 1.

In Fig. 2, we show the ratio of the shock emissivity to the total emissivity (including averaged line processes and Thomson scattering), evaluated at the outer boundary of the wind (solid) and at 1.2 R_* (dash-dotted), corresponding to the onset of X-ray emission in this model. There are a number of interesting features visible:

(i) The total emissivity in the outer wind is dominated by shock emission from just shortward of the He II edge until 2.5 keV (the highest energy we consider in our models). The emissivity in the lower wind, however, is dominated by shock emission only until 200 eV, whilst for larger energies the (local) shock contribution decreases drastically, because the assumed shock temperatures $(\propto (v(r)/v_{\infty})^2)$ are rather low here (≤ 100 kK). The question is then: Which processes dominate the *total* emissivity at high energies in the lower wind? Indeed, this is the re-emission from electron-scattering, being proportional to the mean intensity, and being quite high due to the large number of incoming photons from *above*, i.e., from regions where the shock temperatures are high! This effect becomes also visible in the local radiative fluxes at these frequencies, which are negative, i.e., directed inwards.

(ii) Both in the outer and inner wind, the shock emission is also significant longward from the He II edge, until $\lambda \approx 350$ Å, thus influencing the ionization balance of important ions. Whilst the fluxes of models without shock emission and those with $R_{\min} \gtrsim 2R_*$ display a significant absorption edge for C III and N III (see Fig. 1), these edges have almost vanished in the models with $R_{\min} = 1.2 \dots 1.5 R_*$, because of the dominant shock emissivity increasing the degree of ionization. Even more, all models display fluxes in this region which lie well above those from models without shock emission, because of the higher radiation temperatures compared to the cool wind alone.

(iii) Beyond 350 Å, the shock emissivity becomes almost irrelevant (below 10%), so that the corresponding fluxes are barely affected.

(iv) For the two models with $R_{\min} = 1.2$ and $1.5 R_*$, a prominent emission feature between roughly 300 and 320 Å is visible in Fig. 1. A comparison with Fig. 2 (note the box) shows that this

¹⁷ As shown by Pauldrach et al. (1994), the total shock emissivity is roughly a factor of 50 larger than the corresponding hot plasma free-free emission from hydrogen and helium.

¹⁸ Note that particularly the shock temperature is quite high for such a stellar model, but chosen deliberately to allow for somewhat extreme effects.



Fig. 3. Emergent Eddington fluxes for model S30, with $T_s^{\infty} = 3 \cdot 10^6$ K and $R_{\min}^{\text{input}} = 1.5 R_*$, for different values of f_X , and for a model with an unshocked wind. The histogram-like flux-distribution at highest energies results from our resampling of X-ray emissivities (see Sect. 2.1).

emission is due to dominating shock emission of the lower wind, increasing the temperatures of the radiation field beyond those of the unshocked wind.

Coming back to Fig. 1, significant flux differences between the shocked and the unshocked models are visible for all values of R_{\min} (even for $R_{\min} = 2$ or 10 R_*) below $\lambda \leq 350$ Å, particularly below the N III and C III edges, because of the higher ionization.

On the other hand, the models with $R_{\min} = 1.2$ and $1.5 R_*$ are almost indistinguishable, at least regarding the pseudocontinuum fluxes. This turns out to be true also for He II 1640 and He II 4686, though these lines become sensitive to the choice of R_{\min} if we change R_{\min} from 1.5 to 2 R_* , due to the different intensities around the He II edge and around He II 303 (Lymanalpha) in the line-forming region. We will come back to this point in Sect. 5.1.2.

Impact of f_X . In Fig. 3, we investigate the impact of f_X , which has a most direct influence on the strength of the X-ray emission (cf. Eqs. 1 and 3). Having more X-ray photons leads to higher Xray fluxes/luminosities and to less XUV/EUV-absorption from the cool wind, because of higher ionization stages. The latter effect becomes particularly visible for the model with $f_{\rm X} = 0.1$, which was used to check at which level of X-ray emission we start to change the overall ionization stratification. Most importantly, helium (with He II as main ion beyond 1.2 R_* for S30 models with typical values $0.03 \leq f_X \leq 0.05$) becomes more ionized, reaching similar fractions of He II and He III between 2.2 R_* (~0.5 v_{∞}) and 8.7 R_* (~0.8 v_{∞}). And also the main ionization stage of oxygen (which is O IV in S30 models with typical X-ray emission parameters) switches to O v between 1.8 R_* $(\sim 0.4 v_{\infty})$ and $4.0 R_* (\sim 0.7 v_{\infty})$ when f_X is set to 0.1. The change in the ionization of helium (and oxygen) becomes clearly visible in the much weaker He II edge and much higher fluxes in the wavelength range below 228 Å, compared to models with lower $f_{\mathbf{X}}$.



Fig. 4. Emergent Eddington fluxes for model S30, with $f_X = 0.03$ and $R_{\min}^{\text{input}} = 1.5 R_*$, for different values of maximum shock temperature, T_s^{∞} .

Impact of T_s^{∞} . As displayed in Fig. 4 (see also Pauldrach et al. 2001), the change in the maximum shock temperature, T_s^{∞} , becomes mostly visible for the fluxes shortward of ≈ 60 Å (of course, the hard X-ray band is even more affected, but not considered in our models). Though for the highest maximum shock temperature considered here, $T_s^{\infty} = 5 \cdot 10^6$ K (corresponding to $u_{\infty} \approx 590$ km s⁻¹), we significantly increase the population of the higher ionized atomic species, this is still not sufficient to change the main ionization stages present in the wind.

4.2. Scaling relations for L_x

From analytical considerations, Owocki & Cohen (1999) showed that for a constant volume filling factor (and neglecting effects of radiative cooling, see also below) the optically thin¹⁹ wind X-ray luminosity depends on the square of the mass-loss rate, $L_x \propto (\dot{M}/v_{\infty})^2$, whilst the X-ray luminosity of optically thick winds scales linearly with the mass-loss rate, $L_x \propto \dot{M}/v_{\infty}$, provided that one compares models with the same shock temperatures and assumes a spatially constant X-ray filling factor. These relations become somewhat modified if there is a dependence of T_s on the wind terminal velocity, as adopted in our 'standard' X-ray description (see also KK09).

Note, however, that in a more recent study, Owocki et al. (2013) derived, again from analytic considerations, scaling relations for L_x for radiative and adiabatic shocks embedded in a cool wind. At first glance, their assumptions seem quite similar to those adopted by Feldmeier et al. (1997a) (which is the basis of our treatment), but in the end they predict different scaling relations for *radiative* shocks than resulting from the modeling here. This discrepancy might lead to somewhat different scaling relations for L_x , and needs to be investigated in forthcoming work; for now, we simply compare our models to the earlier results by Owocki & Cohen (1999) (a similar test was done by KK09).

To this end, we calculated S30, S40 and S50 wind models with a fixed X-ray description: $f_X = 0.025$, $m_x = 20$, and $\gamma_x = 0.5$. For our tests we used, for all models, a constant maximum jump velocity, $u_{\infty} = 400$ km s⁻¹ (corresponding to maximum shock

¹⁹ with respect to the cool wind absorption



Fig. 5. Emergent X-ray luminosities (in erg s⁻¹) as a function of \dot{M}/v_{∞} . Supergiant models S30 (asterisks), S40 (triangles) and S50 (squares) with $T_{\rm eff}$ = 30, 40 and 50 kK, respectively, and mass-loss rates between 10⁻⁹ and $2 \cdot 10^{-5} M_{\odot}/{\rm yr}$. All models have the same X-ray properties, $f_x = 0.025$, $\gamma_x = 0.5$, $m_x = 20$, and a maximum jump-velocity, $u_{\infty} = 400$ km s⁻¹, corresponding to maximum shock temperatures of $2.3 \cdot 10^6$ K. The X-ray luminosities have been calculated in the range 0.1 - 2.5 keV (black, green, turquoise), and in the range 0.35 to 2.5 keV (blue, red and magenta). The dashed lines (no fits) serve as guidelines to check the predicted behavior for optically thin (red and green) and optically thick (black) conditions. Note the strong deviation of models S50 (turquoise squares) from the predicted optically thin scaling, when integrating until 100 eV, due to 'normal' stellar/wind radiation just in this energy range. (See text.)

temperatures of $2.3 \cdot 10^6$ K), in order to be consistent with the above assumptions.

For these models (with parameters, except for \dot{M} , provided in Table 1), we varied the mass-loss rates in an interval between 10^{-9} and $2 \cdot 10^{-5} M_{\odot}$ /yr. and integrated the resulting (soft) X-ray luminosities in two different ranges: 0.1 to 2.5 keV and 0.35 to 2.5 keV.

From $\dot{M} \gtrsim 10^{-7} M_{\odot} \text{yr}^{-1}$ on, the wind becomes successively optically thick at higher and higher energies (though, e.g., for $\dot{M} = 10^{-6} M_{\odot} \text{yr}^{-1}$ it is still optically thin below ~ 10 Å, i.e., above 1.24 keV). Indeed, the X-ray luminosities of our corresponding models are linearly dependent on (\dot{M}/v_{∞}) , as can be seen in Fig. 5 by comparing with the black dashed line. For lower \dot{M} , the wind is optically thin at most high energy frequencies, and also here our results follow closely the predictions $(L_x \propto (\dot{M}/v_{\infty})^2)$, when comparing with the red or green dashed lines.

A second finding of Fig. 5 relates to the optically thin scaling for model S50, when either starting the integration at 100 eV (turquoise squares) or at 350 eV (red squares). Whilst for S30 (asterisks) and S40 (triangles) the X-ray luminosities just increase by roughly one dex when including the range from 100 to 350 eV but still follow the predicted scaling relation, the S50 models show an increase of four orders of magnitude for the lowest \dot{M}/v_{∞} values in this situation (and do *not* follow the predictions).

To clarify this effect, Fig. 6 shows the scaled (scaling proportional to R_*^2 and v_{∞}^2) Eddington flux as a function of wavelength and energy, for the supergiant models S30 (black), S40 (green) and S50 (turquoise) with identical, low mass-loss rates, $10^{-8}M_{\odot}$ /yr. Additionally, energies of 100, 150 and 350 eV have



Fig. 6. Logarithmic, scaled Eddington flux (in units of erg cm⁻² s⁻¹ Hz⁻¹) as a function of wavelength/energy, for the supergiant models S30 (black), S40 (green) and S50 (turquoise) with identical mass-loss rates, $10^{-8} M_{\odot}/\text{yr}$. All models have the same X-ray properties, as denoted in Fig. 5. The Eddington fluxes have been scaled by $(R_*/R_{\odot})^2$ and $(v_{\infty}/1000 \text{ km s}^{-1})^2$, to ensure *theoretically* similar values of optically thin X-ray emission. The dotted lines denote energies of 350, 150, and 100 eV, corresponding to 35, 83, and 124 Å. (See text.)

been marked by dotted vertical lines. Beyond 150 eV, all models, independent of their specific parameters, display the same scaled fluxes, thus verifying the optically thin scaling of X-ray luminosities (in this case, only with respect to v_{∞}). For the S50 model, however, the energy range below 150 eV (and, for other parameter-sets, also below even higher energies) is contaminated by 'normal' stellar/wind radiation (which increases as function of T_{eff} ; see also Macfarlane et al. 1994, their Fig. 5), leading to the strong deviation from the optically thin X-ray scaling law as visible in Fig. 5. In so far, the *total* X-ray luminosity (regarding the wind emission) of hotter objects might be overestimated when integrating until 100 eV.

In summary, we conclude that our implementation follows the predicted scaling relations, but we also suggest to choose a lower (in energy) integration limit of 0.15 keV (or even 0.3 keV, to be on the safe side) when comparing the X-ray luminosities of different stars (both with respect to models *and* observations).

In this context, we note that there is a clear distinction between the *observable* soft X-ray and the longer-wavelength, soft X-ray and XUV/EUV emission that is almost never directly observed, but – as already outlined – is very important for photoionizing relevant ions. 'Modern' X-ray observatories such as XMM-NEWTON/RGS and CHANDRA/HETG do not have a response below 0.35 keV and 0.4 keV, respectively²⁰, and even a modest ISM column makes it functionally impossible to see Xray emission below 0.5 keV.

4.3. Comparison with WM-BASIC models

Finally, we checked also the quantitative aspect of our results, by comparing with analogous²¹ WM-BASIC models. As already

²⁰ though ROSAT observed down to 0.1 keV, and also EUVE made a few important measurements relevant for massive stars, in particular for ϵ CMa (B2II), e.g., Cassinelli et al. (1995)

²¹ remember the difference in the velocity fields

Table 2. Left part: X-ray emission parameters used to compare FASTWIND and WM-BASIC models ($u_{\infty}/v_{\infty} = 0.3$ and $\gamma_x = 1.0$). For stellar and wind parameters see Table 1. Right part: L_x/L_{bol} (logarithmic) provided as input for WM-BASIC (WMB), compared with the corresponding output value from FASTWIND (FW), integrated in the frequency range between 0.1 to 2.5 keV. See Sect. 4.3.

Model	$f_{\rm X}$	R_{\min}	u_{∞}	T_s^{∞}	$L_{\rm x}/L_{\rm bol}$	$L_{\rm x}/L_{\rm bol}$				
	(%)	(R_*)	$({\rm km}{\rm s}^{-1})$	(10 ⁶ K)	(WMB)	(FW)				
Dwarfs										
D30	2.00	1.24	532	3.90	-9.4	-9.4				
D35	0.96	1.29	622	5.27	-8.3	-8.5				
D40	1.44	1.21	715	6.98	-7.0	-7.0				
D45	1.38	1.20	894	10.9	-6.4	-6.5				
D50	2.11	1.22	950	12.4	-5.6	-5.8				
Supergiants										
S30	1.99	1.50	453	2.93	-6.3	-6.4				
S35	1.24	1.43	577	4.54	-6.2	-6.3				
S40	0.80	1.33	663	6.00	-6.3	-6.5				
S45	0.93	1.25	754	7.76	-6.2	-6.3				
S50	3.13	1.26	941	12.1	-5.2	-5.4				

pointed out, the X-ray description in both codes is quite similar, and there is only one major difference. In WM-BASIC, the user has to specify a certain value for L_x/L_{Bol} (e.g., 10^{-7} as a prototypical value), and the code determines iteratively the corresponding f_X , whilst the latter parameter is a direct input parameter in the updated version of FASTWIND. In both cases, we used a frequency range between 0.1 to 2.5 keV.

Thus, we first calculated WM-BASIC models with stellar/wind parameters from Table 1, and with X-ray emission parameters from Table 2. For the maximum jump velocity we assumed, as an extreme value, $u_{\infty}/v_{\infty} = 0.3$, together with X-ray luminosities as displayed in the sixth column of Table 2. These values then correspond to the f_X values provided in the second column of the same table, acquired from the WM-BASIC output. We note here that the input values of L_x/L_{Bol} (to WM-BASIC) were not chosen on physical grounds, but were estimated in such a way as to result in similar values for f_X , in the range between 0.01 to 0.03.

To check the overall consistency, we calculated a similar set of FASTWIND models, now using the f_X values from Table 2 as *input*. In case of consistent models, the resulting L_x values (from the output) should be the same as the corresponding input values used for WM-BASIC. Both these values are compared in the last two columns of Table 2. Obviously, the agreement is quite good (not only for the supergiants, but also for the dwarfs), with differences ranging from 0.0 to 0.2 dex, and an average deviation of 0.13 dex.

In a second step, we compared the fluxes resulting from this procedure in Fig. 7. For clarity, the fluxes were shifted by -3, -6, -9, and -18 dex (S35, S40, S45, S50), where the solid lines correspond to the FASTWIND and the dashed lines to the WM-BASIC results.

The comparison shows a remarkably good agreement, with no striking differences. Smaller differences in the lower wavelength range ($\lambda < 100$ Å) are related to a different frequency sampling (without an effect on the total X-ray luminosity). At longer wavelengths, these differences are related to the fact that WM-BASIC provides high-resolution fluxes, whilst FASTWIND calculates fluxes using averaged line-opacities²². Most impor-



Fig. 7. Logarithmic Eddington fluxes as a function of wavelength for supergiant models (see Table 1 and Table 2). The solid lines refer to results from our updated version of FASTWIND, and the dashed ones to WM-BASIC results (Pauldrach et al. 1994, 2001). For clarity, the S35, S40, S45, and S50 model fluxes have been shifted by -3, -6, -9, and -18 dex, respectively.

tant, however, is our finding that the fluxes are not only similar at high frequencies (indicating similar emissivities and cool-wind opacities), but also longward from the He π edge, indicating a similar ionization equilibrium (modified in the same way by the emission from shocked material).

At this stage, we conclude that our implementation provides results that are in excellent agreement with the alternative code WM-BASIC, both with respect to integrated fluxes as well as frequency edges, which moreover follow the predicted scaling relations. Having thus verified our implementation, we will now examine important effects of the X-ray radiation within the stellar wind.

5. Results

In this section, we discuss the major results of our model calculations. In particular, we study the impact of X-ray emission on the ionization balance of important elements, both with respect to direct (i.e., affecting the valence electrons) and Auger ionization. We also discuss the impact of dielectronic recombination and investigate the radial behavior of the high-energy mass absorption coefficient, an essential issue with respect to the analysis of X-ray line emission.

Note that all following results refer to our specific choice of the run of the shock temperature (see Eqs. 6 and 7), which, in combination with our grid-parameter $\gamma_x = 1$, leads to shock temperatures of $T_s(v_{\infty}/2) = 0.25 T_s^{\infty}$ in the intermediate wind at $v(r) = 0.5 v_{\infty}$.

²² for details, see Puls et al. 2005



Fig. 8. Ionization fractions of important ions at $v(r) = 0.5 v_{\infty}$, as a function of T_{eff} , for models with typical X-ray emission (triangles, $f_X = 0.03$, $T_s^{\infty} = 3 \cdot 10^6$ K, corresponding to $u_{\infty} = 460$ km s⁻¹), and without X-rays (asterisks). The solid lines refer to supergiant models, and the dashed ones to dwarf models. For clarity, the ionization fractions of dwarf models have been shifted by one dex. For stellar parameters and onset radius, R_{\min} , see Table 1.

5.1. Ionization fractions

5.1.1. General effects

Though only indirectly observable (particularly via UV resonance lines), ionization fractions provide useful insight into the various radiative processes in the atmosphere. In the following, we compare, for important ions (i.e., for ions with meaningful wind lines), the changes due to the *combined* effects of direct and Auger ionization, whilst the specific effects of Auger ionization will be discussed in Sect. 5.2. These comparisons will be performed for our supergiant (solid) and dwarf models (dashed) from Table 1, and for the center values of our X-ray emission parameter grid (Sect. 3), $f_X = 0.03$, $T_s^{\infty} = 3 \cdot 10^6$ K, that are prototypical in many cases.²³ Comments on the reaction due to different parameters will be given in the next section. All ionization fractions have been evaluated at a representative velocity, $v(r) = 0.5 v_{\infty}$, and are displayed in Fig. 8. To check the influence of X-ray emission, one simply needs to compare the triangles (with) and the asterisks (without X-ray emission).

Carbon. Though our model atom for carbon will be improved soon, already the present one (from the WM-BASIC data base) is certainly sufficient to study the impact of shock radiation. The upper panels of Fig. 8 display the results, which indicate an effect only for 'cooler' supergiant models, with $T_{\rm eff} < 40$ kK. For these objects, C III and C IV become somewhat depleted (less than a factor of ten), whilst C v (which is, without X-ray emission, a trace ion at 30 kK) becomes significantly enhanced. For dwarfs in this temperature range, only C v becomes increased, since the emission (scaling with ρ^2) is still too weak to affect the major ions.²⁴ For models with $T_{\rm eff} > 40$ kK, on the other hand, the temperature is already hot enough that the ionization balance is dominated by the 'normal' stellar radiation field, and no effect due to X-ray emission is visible.

Nitrogen (2nd row) and oxygen (third row of Fig. 8) suffer most from the inclusion of shock radiation. In the following, we concentrate on the differences produced by X-ray ionization in general, whilst in subsequent sections we will consider specific effects.

²³ Note that such maximum shock temperatures might be too high for models around T_{eff} = 30 kK, and that certain effects (as discussed in the following) might thus be overestimated in this temperature range.

 $^{^{24}}$ But note that the actual filling factor in dwarfs might be *much larger* than 0.03, e.g., Cassinelli et al. (1994), Cohen et al. (1997, 2008), Huenemoerder et al. (2012).

Nitrogen. In the 'cool' range, the behavior of N III, N IV and N V is very similar to the corresponding carbon ions (i.e., a moderate depletion of N III and N IV, and a significant increase of N V, particularly at $T_{\rm eff}$ between 30 and 35 K), whereas in the hot range it is different. Here, N III and N IV continue to become depleted, but N v increases only as long as $T_{\rm eff} < 45$ kK, and decreases again at 45 and 50 kK. In other words, when N v is already the main ion for non-X-ray models, it becomes (slightly) depleted when the X-rays are switched on, in contrast to C v which remains unmodified beyond 40 kK. This difference, of course, relates to the fact that C v has a stable noble-gas (He-) configuration, with a high-lying ionization edge (31.6 Å), compared to the N v edge at roughly 126 Å that allows for a more efficient, direct ionization by emission from the shock-heated plasma.

Oxygen. For almost every temperature considered in our grid, the inclusion of X-rays has a dramatic effect on the ionization of oxygen. At 30 kK, O IV becomes the dominant ion²⁵, when for non-X-ray models the main ionization stage is still O III, whereas at the hot end O IV becomes somewhat depleted. The behavior of O v is similar to N v (though the final depletion is marginal), and O vI displays, at all temperatures, the largest effect. At cool temperatures, the ionization fraction changes by 15 orders of magnitude, but even at the hottest $T_{\rm eff}$ there is still an increase by three to four dex. As is well known, this has a dramatic impact on the corresponding resonance doublet.

Silicon. In almost all hot stars, the dominant ion of silicon is Si v (again a noble-gas configuration), and Si IV forms by recombination, giving rise to the well-known Si IV luminosity/mass-loss effect (Walborn & Panek 1984, Pauldrach et al. 1990). The bottom left panel of Fig. 8 displays an analogous dependence. Whilst for dwarfs (low ρ^2) no X-ray effects are visible for Si IV, this ion becomes depleted for cool supergiants ($T_{\rm eff} \leq 35$ kK), at most by a factor of ten.

Phosphorus. During recent years, it turned out that the observed P v doublet at λ 1118,1128 is key²⁶ for deriving mass-loss rates from hot star winds, in parallel with constraining their inhomogeneous structure (Fullerton et al. 2006, Oskinova et al. 2007, Sundqvist et al. 2011, Šurlan et al. 2013, Sundqvist et al. 2014). Thus, it is of prime importance to investigate its dependence on X-rays, since a strong dependence would contaminate any quantitative result by an additional ambiguity.

As already found in previous studies (e.g., KK09, Bouret et al. 2012), also our results indicate that P v is not strongly modified by X-ray emission (middle and right lower panels of Fig. 8), though more extreme X-ray emission parameters, e.g., $f_X = 0.05$ and/or $T_s^{\infty} = 5 \cdot 10^6$ K, can change the situation (see section 5.1.3). Even more, the apparently small change in the ionization fraction of P v at typical X-ray emission parameters (decrease by a factor of two to three) can still be of significance, given the present discussion on the precision of derived mass-loss rates (with similar uncertainties).

Regarding the ionization of P vI, cold models (30 and 35 kK) change drastically when X-ray emission has been included, both for supergiants and dwarfs. Since we find less P vI in hot models with shocks (compared to models without), this indicates that the ionization balance is shifted towards even higher stages (P vII).

In this context, we note that Krtička & Kubát ($\overline{2012}$) investigated the reaction of P v when incorporating additional, strong XUV emissivity (between 100 and 228 Å) and micro-clumping



Fig. 9. Helium ionization fractions as a function of local velocity, for an S30 model with ($f_{\rm X} = 0.03$ and $T_{\rm s}^{\infty} = 3 \cdot 10^6$ K) and without X-rays. See text.

into their models. The former test was driven by a previous study by Waldron & Cassinelli (2010) who argued that specific, strong emission lines in this wavelength range could be of significant impact. Indeed, Krtička & Kubát (2012) were able to confirm that under such conditions²⁷ P v becomes strongly depleted, in parallel with changes in the ionization fractions of, e.g., C Iv, N IV, and O IV (see also Sect. 5.1.3). Further work is certainly required to identify the source of such additional emissivity (if present), and, in case, incorporate this mechanism into our models.

5.1.2. Impact on helium

During our analysis, we noted that also helium can be affected by shock emission (see also Sect. 4.1), a finding that has been rarely discussed in related literature. In particular, He II (and He I) can become depleted in the intermediate wind, though only for our 'cooler' supergiant models with 30 kK $\leq T_{\text{eff}} \leq 40$ kK. The effect is strongest for S30 models, but barely noticeable already at S40, independent of the specific X-ray emission parameters. For all our dwarf models, no changes are visible at all.

Figure 9 displays the helium ionization fractions for an S30 model with typical X-ray emission parameters, as a function of local velocity. The depletion of He II (and, in parallel, of He I that is not displayed) is significant in the region between $0.2v_{\infty} \leq v(r) \leq 0.8v_{\infty}$, and results from the increased ionization due to the increased radiation field (in the He II Lyman continuum) in models with shocks (note also the corresponding increase of He III).

In Fig. 10, we compare the helium ionization fractions from our solution and a corresponding WM-BASIC S30 model, but now with X-ray emission parameters as tabulated in Table 2 (the major difference is a filling factor of 0.02 instead of 0.03). Here, we display the fractions as a function of τ_{Ross} , to enable a comparison of the photospheric regions as well. Again, the depletion of He II (now located between $\tau_{Ross} \approx 0.1...0.01$) is visible, and our results coincide perfectly with those predicted by WM-BASIC.

²⁵ this is also true for models with different X-ray emission parameters ²⁶ because it is the only UV resonance line(-complex) that basically never saturates, due to the low phosphorus abundance

²⁷ enhanced emissivity in the XUV range; note, however, that the lines refered to by Waldron & Cassinelli (2010) are included in 'standard' plasma emission codes



Fig. 10. Helium ionization fractions as a function of τ_{Ross} , for S30 models calculated by FASTWIND and WM-BASIC, both with X-ray emission parameters from Table 2. The agreement is excellent.

Since the ionization balance changes already at quite low velocities, this might affect at least two important strategic lines, He II 1640 and He II 4686.²⁸ From Fig. 11, we see that this is actually the case: He II 4686 displays stronger emission, whilst He II 1640 displays a stronger emission in parallel with absorption at higher velocities, compared to the non-X-ray model (dotted). This is readily understood since He II 4686 is predominantly a recombination line, such that the increase in He III leads to more emission; to a lesser extent, this is also true for He II 1640. The lower level of this line, n = 2 (responsible for the absorption), is primarily fed by pumping from the ground-state via He II 303. We have convinced ourselves that the increased pumping because of the strong EUV radiation field leads to a stronger population of the n = 2 state (even if He II itself is depleted), so that also the increased absorption is explained.

As already pointed out in Sect. 4.1, changing R_{\min} from 1.5 to 1.2 R_* does not make a big difference. Increasing R_{\min} to 2 R_* , however, changes a lot, as visible from the dash-dotted profiles in Fig. 11. Except for slightly more emission (again because of increased He III in regions with $r > 2R_*$), the difference to profiles from models without shock emission becomes insignificant, simply because both lines predominantly form below the onset radius.

5.1.3. Dependence on filling factor and shock temperature

As we have seen already above, each ion reacts somewhat differently to the imposed shock radiation. In this section we describe how a change of important X-ray characteristics affects important ions. The figures related to this section are enclosed in Appendix A. The upper figure on each page shows specific ionization fractions with and without X-rays, as a function of $T_{\rm eff}$, for our supergiant and dwarf models (S30 to S50 and D30 to D50, respectively). The ionization fractions have been evaluated at the location where the impact of shock radiation is most evident for the considered ion. Each of these figures contains nine panels, where both the filling factor and the maximum shock temperature are varied according to our grid, i.e., $f_{\rm X} = 0.1, 0.3$,

Fig. 11. Synthetic He II 1640 and He II 4686 profiles for our S30 model. Each profile corresponds to a different X-ray description. Solid: $f_X = 0.03$, $T_s^{\infty} = 3 \cdot 10^6$ K, $R_{\min} = 1.5$ R_* ; dash-dotted: as solid, but with $R_{\min} = 2$ R_* ; dotted: no shock emission.

0.5 and $T_s^{\infty} = 1,3,5\cdot10^6$ K. Note that the onset radius, R_{\min} , was set to its default value for all models. The lower two figures on each page display the ionization fractions for our dwarf (left) and supergiant models (right), evaluated at the same location as above, but now overplotted for all values of f_X (different colors) and T_s^{∞} (different symbols), and without a comparison to the non-X-ray case. Thus, the upper figure allows to evaluate the X-ray effects in comparison to models without shock emission, whilst the lower two figures provide an impression on the differential effect, i.e., the range of variation.

Carbon. C III and C IV are significantly affected in supergiant models with 30 kK $\lesssim T_{\text{eff}} \lesssim 40$ kK, for intermediate to large values of f_X and T_s^{∞} . The depletion of C III and C IV reaches a factor of 10 (or even more) in cooler supergiant models when the highest values of X-ray emission parameters are adopted, which is reflected in a corresponding increase of C v. On the other hand, C III and C IV are barely modified in supergiant models with the lowest values of f_X or T_s^{∞} , which is also true for dwarf models with any value of our parameter grid (see Figs. A.1/A.2). The ionization fraction of C v increases also for the lowest values of X-ray emission parameters, once more for the cooler models (here, also dwarfs are affected). C v remains unmodified beyond 40 kK due to its stable noble-gas configuration, as previously noted.

Nitrogen. The behavior of N III, N IV and N V in the colder models is similar to the corresponding carbon ions, for all different X-ray descriptions. For higher T_{eff} , increasing f_X enhances the depletion of N III and N IV in both supergiants and dwarfs, whilst the impact of T_s^{∞} is rather weak. At the largest values of X-ray emission parameters, both stages become highly depleted (one to two orders of magnitude) for all models but D30 and D35.

Shock radiation is essential for the description of N v at almost any temperature, particularly for models with $T_{\rm eff} < 45$ kK (Figs. A.3/A.4). Here, the increase of N v (compared to non-X-ray models) can reach 4 to 5 dex at the lowest temperatures. At 45 kK, only a weak impact of shock radiation can be noted, whilst for 50 kK a high depletion of N v for extreme parameters values becomes obvious. Once more, the impact of f_X is more prominent than of T_s^{∞} , mainly for the coldest models where N v becomes enhanced by one order of magnitude when increas-

 $^{^{\ 28}}$ Most other He II and He I lines are formed in the photosphere, and remain undisturbed.

ing f_X from 0.01 to 0.05 and keeping T_s^{∞} constant. The hottest models with moderate to high parameters ($f_X \gtrsim 0.02$ and $T_s^{\infty} \gtrsim 2 \cdot 10^6$ K) indicate that also N vI becomes strongly affected by changes in the X-ray ionization.

Oxygen. Independent of the X-rays description, the depletion of O tv for hot models happens only in a specific range of the wind, between 0.4 to 0.8 v_{∞} (similar to the case of He II discussed in the previous section). Also for X-ray emission parameters different from the central value of the grid, the behavior of O v is still very similar to N v, where mainly the cold models are quite sensitive to variations of f_X (Figs. A.5/A.6). The shock radiation increases the ionization fraction of O v by 5 to 6 dex (when f_X varies between 0.01 and 0.05, independent of T_s^{∞}) for the coolest models, whilst these factors decrease as T_{eff} approaches 40 to 45 kK. Models with $T_{\text{eff}} = 45$ kK are barely affected, independent of the specific X-ray emission parameters. Similar to the case for N v at highest values of f_X , T_s^{∞} , and T_{eff} , the corresponding depletion of O v points to the presence of a significant fraction of higher ionization stages.

As pointed out already in Sect. 5.1.1 (see also Sect. 5.2), the X-ray radiation is essential for the description of O vI, which shows, particularly in the cold models, a high sensitivity to both f_X and T_s^{∞} (Figs. A.7/A.8).

Silicon. Also when varying the X-rays description, Si IV still remains unaffected from shock emission in dwarf models. On the other hand, for cool supergiants ($T_{\text{eff}} \leq 35$ kK), Si IV becomes even more depleted when f_{X} increases (though T_{s}^{∞} has a negligible influence). No variation is seen in Si v, as expected due to its noble-gas configuration.

Phosphorus. Pv shows a sensitivity to both f_X and T_s^{∞} , but in this case T_s^{∞} is more relevant. Though no difference between models with and without shocks is seen for the lowest values of T_s^{∞} , particularly the supergiant models develop a depletion with increasing shock temperature, even at lowest f_X . As noted already in Sect. 5.1.1, for extreme X-ray emission parameters the depletion of Pv is significant for all models (both supergiants and dwarfs), except for D30 (Figs. A.9/A.10). Finally, even Pv becomes highly depleted for hot models ($T_{\text{eff}} \ge 40 \text{ kK}$) at intermediate and high values of T_s^{∞} , which indicates the presence of even higher ionization stages.

To summarize our findings: When increasing the values for f_X and T_s^{∞} , the effects already seen in Fig. 8 become even more pronounced, as to be expected. For most ions, the impact of f_X appears to be stronger than the choice of a specific T_s^{∞} (provided the latter is still in the range considered here), though P v and O vI (for the cooler models) show quite a strong reaction to variations of the latter parameter. Overall, the maximum variation of the ionization fractions within our grid reaches a factor of 10 to 100 (dependent on the specific ion), where lower stages (e.g., C IV, N IV, O IV and P V) become decreased when f_X and T_s^{∞} are increased, whilst the higher stages (e.g., N v, O v, O vI) increase in parallel with the X-ray emission parameters. Only for Si IV, the impact of X-rays remains negligible in all models except for S30 and S35.

5.1.4. Comparison with other studies

Since the most important indirect effect of shock emission is the change in the occupation numbers of the *cool* wind, it is worth-while and necessary to compare the ionization fractions resulting from our implementation with those presented in similar studies.

To this end, (i) we recalculated the models described in KK09, (ii) compared with two models (for HD16691

Fig. 13. Radial stratification of phosphorus ionization fractions, as a function of τ_{Ross} , for our model of HD 203064 at T_{eff} = 34.5 kK (see KK09 for stellar, wind and X-ray emission parameters). In our implementation, P v is barely modified by the X-ray radiation field, whilst a considerable impact is seen for P vI.

Fig. 14. Ionization fractions of P_{IV} (asterisks) and P_V (triangles) as a function of normalized velocity, for an S35 (solid) and S40 (dashed) model. Both models have been calculated with a clumping factor $f_{cl} = 20$, and a mass-loss rate reduced by a factor of ~4 compared to the values provided in Table 1. Compare with Fig. 10 in Bouret et al. (2012).

and HD 163758) presented in Bouret et al. (2012) who used CMFGEN and SEI²⁹-fitting to calculate/derive the ionization fractions of phosphorus, and (iii) compared with the ionization fractions predicted by WM-BASIC.

Regarding the first point, we recalculated the 14 O-star models (in the temperature range between 30 and 40 kK) presented by KK09, using parameters from their Tables 2 and 3, both without and with shock emission ($f_X = 0.02$ and $u_{\infty}/v_{\infty}=0.3$), by means of FASTWIND, using H, He, C, N, O, Si and P as explicit ions. Figure 12 shows our results for the ionization fractions of selected ions, as a function of T_{eff} , and evaluated at $v(r) = 0.5v_{\infty}$. The layout of this figure is similar to Figure 8 in KK09, and has been augmented by O vI evaluated at $v(r) = 0.05v_{\infty}$ and N v evaluated at $v(r) = 0.8v_{\infty}$, corresponding to their Figures 9 and 10.

²⁹ Sobolev with exact integration (Lamers et al. 1987)

Fig. 12. Ionization fractions of selected ions as a function of T_{eff} , for 14 O-star models as detailed in Krtička & Kubát (2009, KK09) and recalculated by us using FASTWIND. If not indicated otherwise, fractions are displayed at $v(r) = 0.5v_{\infty}$. As in previous figures, triangles represent models with shocks, and crosses those without. This figure reproduces, in most parts, the layout of Figure 8 from KK09, such that differences and similarities between our and their results can be easily recognized. For details, see text.

Indeed, there are only few ions which display similar fractions over the *complete* temperature range of the O-star models considered by KK09 (which still misses the hotter O-stars beyond 40 kK). For C IV, an agreement is present only for the coolest regime ($T_{\rm eff} \leq 32$ kK) where both studies predict C IV as the main ion, independent whether X-rays are present or not. Whilst the fractions for non-X-ray models are comparable also for hotter temperatures, the X-ray models by KK09 show a much larger depletion of C IV (fractions of 10^{-2} to 10^{-3} for $T_{\rm eff} >$ 34 kK) than our models (still above 10^{-1}).

For O vI, agreement between both results is present only at the hottest temperatures, whilst between 30 kK < $T_{\rm eff} \leq 37$ kK our models display a factor of ~100 lower fractions, for both the non-X-ray models and the models with shock emission. The same factor is visible in the lower wind ($v(r) = 0.05v_{\infty}$) for the X-ray models, but the non-X-ray models are similar here.

For nitrogen (N IV and N V), on the other hand, the results are quite similar in most cases. The exception is N v for models without shocks, where our results are lower (by ~1 dex) in the intermediate and outer wind ($v(r) = 0.8v_{\infty}$).

For Si IV, both results fairly agree for the X-rays models, though we do not see a significant effect from including the

shock emission in our calculations (in other words, X-ray and non-X-ray models yield more or less identical results). In contrast, the models by KK09 indicate a small depletion of Si rv when including the shock emission, by factors of roughly 2 to 3. Thus, our non-X-ray models have less Si rv than those by KK09.

Again, phosphorus (in particular, Pv) has to be analyzed in more detail. Comparing the last two panels of Fig. 12 with Fig. 8 from KK09, we see that our ionization fractions for Pv agree with KK09 in the coolest models, and in the hottest models regarding Pvi. In the other temperature ranges, however, differences by a typical factor of 2 (regarding Pv) and 2 to 5 (regarding Pvi) are present. In their Fig. 12, KK09 display the radial stratification of the phosphorus ionization fractions for their model of HD 203064, whilst the corresponding results from our implementation are shown in Fig. 13. Both codes yield quite similar fractions for PIV and PV (with and without X-rays) in the external wind. The same is true for Pvi in the model with X-rays, but we have considerably less Pvi for the non-X-ray model. Prominent differences are visible in the lower wind and close to the lower boundary. We attribute this difference to a boundary condition (in the models by KK09) at quite low optical depths, where the electron temperature is still close to the

effective one.³⁰ Thus far, it is conceivable that quite a low ionization stage (P $_{IV}$) dominates their internal atmosphere (followed by P v and negligible P $_{VI}$), whilst in our case it is just the other way round, and P $_{VI}$ dominates, because of the much higher temperatures.

To check these discrepancies further, we compared our results also with calculations performed with CMFGEN. Particularly, we concentrated on two supergiant models at roughly 35 kK and 40 kK (HD 163758 and HD 16691, respectively), as described by Bouret et al. (2012). In these models, an X-ray emitting plasma with *constant* shock-temperature, $T_s(r) =$ $3 \cdot 10^6$ K, a filling factor corresponding to $L_x/L_{bol} = 10^{-7}$, and an onset radius corresponding to 200 to 300 km s⁻¹ was used (J.-C. Bouret, priv. comm.). In Fig. 14, we present our results for P iv and P v which can be compared with their Fig. 10, displaying P v alone. Though our models³¹ do not have identical parameters (in particular, our shock temperatures increase with velocity), the ionization fractions behave quite similar: In the cooler model (solid), the ionization of Pv decreases with velocity, and in the hotter one (dashed), it increases outwards. This is because in the cooler model, Pv is the dominant ion at low velocities, recombining to PIV, whilst in the hotter model PVI dominates at low velocities, recombining to P v in the run of the wind. Of course, there are some quantitative differences, particularly in the intermediate wind³², but we attribute these to a different stratification of the clumping factor, f_{cl} , and to the different description of the X-ray emitting plasma (concerning the reaction of P v on various X-ray emission parameters, see Fig. A.10).

As a final test, we compared our solutions to the predictions by WM-BASIC, using our dwarf and supergiant models (Table 1 and X-ray emission parameters from Table 2). The results are displayed in Figs. B.1 and B.2 (Appendix B). Note that the range of comparison extends now from 30 to 50 kK, i.e., to much hotter temperatures than in the comparison with KK09.

Overall, the agreement between FASTWIND and WM-BASIC is satisfactory, and all trends are reproduced. However, also here we find discrepancies amounting to a factor of 10 in specific cases, particularly for Si IV. Typical differences, however, are on the order of a factor of two or less. We attribute these discrepancies to differences in the atomic models, radiative transfer and the hydrodynamical structure, but conclude that both codes yield rather similar results, maybe except for Si IV which needs to be reinvestigated in future studies.

In Fig. B.3 we see how some of the encountered differences (compared at only *one* depth point, $v(r) = 0.5v_{\infty}$, except for N v) translate to differences in the emergent profiles. As prototypical and important examples, we have calculated line profiles for N v 1720, N v 1238,1242, O v 1371, O v 1031,1037 and P v 1117,1128, and compare them with corresponding WM-BASIC solutions for models S30, D40, S40, D50 and S50 (for model D30, all these line are purely photospheric, and thus not compared). Both the WM-BASIC and the FASTWIND profiles have been calculated with a radially increasing microturbulence, with maximum value $v_{turb}(max) = 0.1v_{\infty}$, which allows for reproducing the blue absorption edge and the 'black trough' (see Sect. 2.1) in case of saturated P Cygni profiles.

This comparison clearly shows that in almost all considered cases the agreement is satisfactory (note that WM-BASIC includes the photospheric 'background', whilst FASTWIND only accounts for the considered line(s)), and that larger differences are present only (i) for N IV and O V in the outer wind, where FASTWIND produces more (N IV) and less (O V) absorption, respectively, and (ii) for strong P v lines, where FASTWIND predicts higher emission.

5.2. Impact of Auger Ionization

All X-ray models discussed so far include the effects from direct *and* Auger ionization, which was shown to play an important role for the ionization balance in stellar winds (e.g., Cassinelli & Olson 1979, Olson & Castor 1981, Macfarlane et al. 1994, Pauldrach et al. 1994). In the following, we investigate the contribution of the latter effect to the total ionization in more detail, particularly since there is still a certain debate on this question.

Figure 15 displays how specific ions are affected throughout the wind, for dwarf and supergiant models with different $T_{\rm eff}$ and typical X-ray emission parameters ($f_{\rm X} = 0.03$ and $T_{\rm s}^{\infty}$ = 3·10⁶ K). Each ion is shown at three different locations: v(r) =0.3 v_{∞} (close to the onset of the shock emission), $v(r) = 0.6 v_{\infty}$ (intermediate wind) and $v(r) = 0.9 v_{\infty}$ (outer wind).

Two general comments: (i) Significant effects are to be expected only for quite high ionization stages, since in the majority of cases Auger ionization couples ions with a charge difference of two (but see Sect. 2.2). E.g., C IV should remain (almost) unmodified, since CII is absent in O- and at least early B-stars, and the K-shell absorption of C_{IV} (with threshold at 35.7 Å), resulting in the formation of C v (charge difference of one!), is in most cases (but see below) negligible compared to the direct ionization of C_{IV} (with threshold at ~192 Å for the groundstate ionization; remember that the radiation field is stronger at longer wavelengths, which favors direct vs. Auger ionization). In contrast, O vi should become significantly affected, since O iv is strongly populated in O-stars, and the transition threshold for the direct ionization from O v (at ~ 109 Å) is now closer to the K-shell edge. Consequently, the transition rates (depending on the corresponding radiation field) are more similar than in the case of C IV.

(ii) In the same spirit, Auger ionization should become negligible, at least in most cases, for the hotter O-stars (see also Sect. 4). Once T_{eff} is high, more direct ionization is present (because of the stronger radiation field at the corresponding, lower-frequency edges), and consequently the impact of Auger ionization should decrease. Though this argumentation is basically correct, the actual results depend, of course, also on the wind-strength, since higher densities lead to more X-ray emission (for identical f_X), which increases the impact of Auger-ionization. E.g., if we check for the behavior of N vi at 0.9 v_{∞} in Fig. 15, we see that for D40, D45 and D50 there is indeed no effect, whilst for S40 and S45 Auger ionization still has a certain influence.

Now to more details. At first note that all ions from C, N, O, Si and P that are *not* displayed in Fig. 15 are barely changed by Auger ionization, with a maximum difference of ± 0.08 dex (corresponding to factors of 0.8 to 1.2) in the fractions calculated with and without Auger.

For carbon, Cv is the only ion which under specific conditions becomes affected by Auger ionization. As visible in the first line of Fig. 15, cold supergiant models display an increase of Cv in the outer wind when Auger is included, since in this case

³⁰ Indeed, we were not able to find statements or figures related to the *photospheric* structure of their models in any of the papers by Krtička and co-workers, so our argument is somewhat speculative.

³¹ S35 and S40, but using a clumped wind with reduced mass-loss rates to ensure comparable wind structures

 $^{^{32}\,}$ J.-C. Bouret provided us with an output of the ionization fractions for P ν and P $\nu.$

Fig. 15. Ionization fractions of ions most affected by Auger ionization, at different depth points. All models have typical X-ray emission parameters ($f_{\rm X} = 0.03$ and $T_{\rm s}^{\infty} = 3 \cdot 10^6$ K). The triangles represent models including Auger ionization (standard approach, similar to Fig. 8), and squares models without (i.e., 'only' direct ionization has been considered). Solid lines refer to supergiant models, and dashed ones to dwarf models. For clarity, the ionization fractions of dwarf models have been shifted by one dex.

the radiation field at the corresponding K-shell edge becomes quite strong, compared to the radiation field around 192 Å (see Fig. 7). This increase is compensated by a similar decrease of C $_{\rm IV}$, which, in absolute numbers, is quite small though.

N vI (second line in Fig. 15) is the only nitrogen ion where larger changes can be noted. In cool dwarfs, it becomes influenced already at 0.3 v_{∞} , and also in the intermediate wind, which is also true for model S30. In the outer wind, differences appear clearly for all models, except for dwarfs with $T_{\rm eff} \gtrsim 40$ kK. The corresponding change in N IV, on the other hand, is marginal, again because N VI itself has a low population, even when Auger is included.

O v behaves similar to N v (mostly no changes), but now a weak effect appears in the outer wind of cool supergiants (third line of Fig. 15), and even for O v1 (compare to the reasoning above), changes in the lower and intermediate wind are barely visible (if at all, then only for the S30 model, see last line of Fig. 15). In the outer wind, however, considerable differences in O v1 (up to three orders of magnitude) can be clearly spotted for all supergiants and cooler dwarf models, similar to the case of N v1. Only for the hottest models, the effect becomes weak. Fig. 16 shows an example for an S40 model where the second-

most populated oxygen ion (O v) changes to O vI after the inclusion of Auger ionization.

Finally, the K-shell edges for phosphorus (not implemented so far) and silicon (with quite low cross-sections) are located at such high energies (> 2 keV or > 6 Å) that the corresponding Auger rates become too low to be of importance, at least for the considered parameter range.

To conclude, in most cases the effects of Auger ionization are only significant in the outer wind (for a different run of shock temperatures, they might become decisive already in the lower or intermediate wind), and for highly ionized species. The effect is essential for the description of N vI and O vI, particularly in the outer wind. Thus, and with respect to strategic UV resonance lines, it plays a decisive role only in the formation of O vI 1031,1037 (but see also Zsargó et al. 2008).

5.3. Dielectronic Recombination of Ov

After comparing the results from our first models accounting for shock-emission with corresponding WM-BASIC results, it turned out that in a specific parameter range (for dwarfs around 45 kK) both codes delivered largely different fluxes around the O IV edge

Fig. 16. Radial stratification of oxygen ionization fractions, as a function of τ_{Ross} , for an S40 model with $f_X = 0.03$ and $T_s^{\infty} = 3 \cdot 10^6$ K. Auger ionization notably affects the presence of O vI in the outer wind ($\tau_{\text{Ross}} \le 10^{-2}$ corresponding to $r \ge 4 R_*$ or $v(r) \ge 0.7 v_{\infty}$). The model without Auger ionization has more O v than O vI, and vice versa when the effect is included.

Fig. 17. Ionization fractions of oxygen, as a function of τ_{Ross} , for a D45 model with $f_{\text{X}} = 0.03$ and $T_{\text{s}}^{\infty} = 3 \cdot 10^6$ K, with and without dielectronic recombination (DR). Note the large differences for all the stages when $\tau_{\text{Ross}} \leq 10^{-2} (v(r) \geq 0.05v_{\infty})$, particularly the change in the main ionization stage (from O v/O vI to O Iv) when DR is included.

at ~160 Å, which could be tracked down to completely different ionization fractions of oxygen. In particular, our models displayed more O v and less O IV than calculated by WM-BASIC.

After investigating the origin of this discrepancy, it turned out that we, inadvertently, had not included the data for dielectronic recombination³³ (hereafter DR) in our oxygen atomic model.³⁴ Thus, DR processes had not been considered for oxygen. A series of studies had recently reconsidered the effects of DR with respect to *nitrogen* (Rivero González et al. 2011, 2012a,b), however no significant effects were found, particularly concerning the formation of the prominent N III $\lambda\lambda$ 4634-4640-4642 emission lines that was previously attributed to DR processes (Bruccato & Mihalas 1971, Mihalas & Hummer 1973).

Nevertheless, we subsequently included DR also into our oxygen atomic model, and were quite surprised by the consequences. Though in a large region of our model grid the changes turned out to be negligible for the fluxes, in all supergiant models and in the dwarf models around 45 kK the ionization fractions were strongly affected, leading to a decrease of O v, typically by a factor of 10 to 50.

For our most problematic D45 model, DR proved to be essential to even predict the correct main ion throughout the wind, and to produce a reliable SED around the O rv edge. Fig. 17 displays the impact of DR for this model. Indeed, the population of every ionization stage becomes modified in the wind, but for O rv this difference is large enough to change it to the main stage of the model. The reason for such drastic impact in the region around D45 is based on the fact that only here the X-ray ionization is potentially able to allow for the dominance of O v (see Fig. 8), which then can be compensated by quite strong dielectronic recombination rates.³⁵

Nevertheless, since in the majority of models O v becomes severely depleted (see above), independent of whether it is a main ion or not, and also O vi is affected, this leads to considerable changes in the corresponding UV lines. Thus, we conclude that DR is inevitable for a correct treatment of oxygen. Moreover, because of this strong impact, the precision of corresponding data needs to be re-checked.

As a final remark, let us note that the inclusion of DR has also an impact on non-X-ray models, but to a much lower extent.

5.4. Mass Absorption Coefficient

As already mentioned in Sect. 1, also the X-ray line emission (observed by means of CHANDRA and XMM-NEWTON) has been modeled and analyzed during recent years, by various groups. Such analysis particularly allows us to obtain constraints on the presence, structure, and degree of wind inhomogeneities at X-ray wavelengths (e.g., Oskinova et al. 2006, Sundqvist et al. 2012a, Leutenegger et al. 2013b), to independently 'measure' the mass-loss rates of O-star winds (e.g., Hervé et al. 2013, Cohen et al. 2014b, Rauw et al. 2015), and even to derive nitrogen and oxygen abundances³⁶ (Oskinova et al. 2006, Zhekov & Palla 2007, Nazé et al. 2012, Leutenegger et al. 2013a). One of the assumptions made by various authors is to consider the mass absorption coefficient of the cool wind material, $\kappa_{\nu}(r)$, as spatially constant, which simplifies the analysis (Owocki & Cohen 2006, Leutenegger et al. 2013b, Cohen et al. 2014b). Other groups include detailed predictions for the spatial and frequency dependence of $\kappa_{\nu}(r)$, calculated by means of POWR (e.g., Oskinova et al. 2006) or CMFGEN (e.g., Hervé et al. 2013, Rauw et al. 2015), and there is an ongoing dis-

³³ This process can be summarized as 'the capture of an electron by the target leading to an intermediate doubly excited state that stabilizes by emitting a photon rather than an electron' (Rivero González et al. 2012a).

 $^{^{34}\,}$ For Si, P and C v \rightarrow C v corresponding data are still missing in our database.

³⁵ As an independent check of our findings, we also calculated WM-BASIC models without DR, and they turned out to be consistent with our non-DR models.

³⁶ Primarily, these abundance determinations involve measuring the strengths of corresponding emission lines in the soft X-ray regime, maybe correcting them for absorption. Note, however, that this diagnostics is *not* a wind absorption diagnostics, but that absorption is only a correction needed to derive line luminosities.

cussion whether the assumption of a spatially constant κ_{ν} is justified and in how far it affects the precision of the deduced massloss rates. Though Cohen et al. (2010, 2014b) have investigated the variation of $\kappa_{\nu}(r)$ and its influence on the derived parameters based on selected CMFGEN-models (also accounting for variations in the CNO-abundances), a systematic study has not been performed so far, and in this section we will do so.

At first, let us consider why and under which conditions κ_{ν} should become more or less spatially constant. The prime reason for this expectation is the fact that the K-shell cross sections (at threshold and with respect to wavelength dependence) of the various ions of a specific atom are quite similar, and that the corresponding edges (for these ions) lie close together. Provided now that (i) *all* ions which are present in the wind are actually able to absorb via K-shell processes, and (ii) that there are no 'background' opacities from other elements, $\kappa_{\nu}(r)$ indeed becomes (almost) spatially constant, since the total opacity is then the simple sum over the K-shell opacities from all contributing atoms,

$$\kappa_{\nu}(r) \approx \sum_{k} \left(\sum_{j} \frac{n_{k,j}(r)}{\rho(r)} \sigma_{k,j}(\nu) \right) \approx$$
$$\approx \sum_{k} \left(\sum_{j} \frac{n_{k,j}(r)}{\rho(r)} \right) \sigma_{k}(\nu) \approx$$
$$\approx \sum_{k} \frac{n_{k}(r)}{\rho(r)} \sigma_{k}(\nu) \approx \frac{\sum_{k} \alpha_{k} \sigma_{k}(\nu)}{m_{\mathrm{H}}(1+4Y_{\mathrm{He}})} := \kappa_{\nu}^{\mathrm{appr}}, \tag{9}$$

with α_k the elemental abundance, Y_{He} the helium abundance (both quantities normalized to hydrogen), and m_{H} the hydrogen mass. k denotes the atomic species, and j the ion, $n_{k,j}$ is the occupation number of ion (k,j), and $\sigma_{k,j} \approx \sigma_k$ the K-shell cross section, being almost independent of j. In the last step of the above derivation, we have assumed that the atmosphere consists mostly of hydrogen and helium.

Thus, we have to check under which conditions restrictions (i) and (ii) might no longer be valid. For the light and abundant elements CNO, K-shell absorption is no longer possible for C v, N vI and O vII. For these ions, only 'ordinary', outer-shell ionization is present, but also here the cross-sections are not too different from the K-shell processes (both with respect to strength and location of edge). Thus, even for highly ionized winds (hot or with strong X-ray emission) where C v, N vI, and O vII are actually present somewhere, the above approximation is still justified. In so far, restriction (i) should play no role, since even higher ionization stages are not too be expected to be significantly populated.

Regarding restriction (ii), the situation is different. The prime background is given by the He II bound-free opacity, which becomes strong in 'cool' and/or helium-recombined winds³⁷, where in the following we always refer to the recombination of He III to He II. Note that already Hillier et al. (1993) showed the importance of outer-wind helium recombination on wind opacity and emergent soft X-ray emission.

Let's check on the maximum influence of the He II boundfree opacity at important K-shell edges. For a crude estimate, we approximate its frequency dependence by $(\nu_0/\nu)^3 = (\lambda/\lambda_0)^3$, and assume the worst case that He II is the only He ion present in the wind. Then, a lower limit for the opacity ratio at specific K-shell

0.68 0.90 1.12 1.34 1.56 1.78 2.00 2.22 2.44 2.66 2.88 $\log (\kappa [\text{cm}^2 \text{g}^{-1}])$

Fig. 18. Contour plots illustrating the radial dependence of the mass absorption coefficient, $\kappa_v(r)$, as a function of wavelength. The upper panel refers to model D30, and the lower to model S40, both with typical X-ray emission parameters ($T_s^{\infty} = 3 \cdot 10^6$ K and $f_X = 0.03$). The positions of the C v edge (outer-shell ionization) and the C rv and O rv K-shell edges are indicated.

edges can be approximated by

$$\frac{\kappa_{k}}{\kappa_{\text{HeII}}}(\lambda_{0}(k)) \approx \frac{n_{k}}{n_{\text{HeII}}} \frac{\sigma_{0}(k)}{\sigma_{0}(\text{HeII})} \left(\frac{\lambda_{0}(\text{HeII})}{\lambda_{0}(k)}\right)^{3} \gtrsim \\ \gtrsim \frac{\alpha_{k}}{\alpha_{\text{He}}} \frac{\sigma_{0}(k)}{\sigma_{0}(\text{HeII})} \left(\frac{228 \text{ Å}}{\lambda_{0}(k)}\right)^{3}, \tag{10}$$

where σ_0 is the cross section at the corresponding edge. Using solar abundances from Asplund et al. (2009), $\lambda_0(C) \approx 35$ Å and $\lambda_0(O) \approx 20$ Å, $\sigma_0 \approx 1.6$, 0.9, and $0.5 \cdot 10^{-18}$ cm² for the threshold cross sections of He II, carbon (K-shell), and oxygen (K-shell), respectively, we find $\kappa_C/\kappa_{\text{HeII}}(35 \text{ Å}) \gtrsim 0.42$ and $\kappa_O/\kappa_{\text{HeII}}(20 \text{ Å}) \gtrsim$ 2.3. Thus, for cool and/or He-recombined winds, the He II opacity dominates at the carbon K-shell edge, whilst at the oxygen edge the K-shell opacities are quite a bit larger than the background. Thus, we would predict that somewhat below \approx 20 Å (beyond 620 eV) restriction (ii) becomes valid, and that κ_v should become depth-independent. Vice versa, the mass absorption coefficient should vary with radius longward from the oxygen or carbon K-shell edge, whenever the background mass absorption coefficient varies, mostly due to changes in the He II ionization throughout the wind.

³⁷ Additionally, the outer-shell ionization of O tv with edge at ≈ 160 Å and the bound-free opacities from other, strongly abundant ions can play a minor role, particularly if He II is weak or absent.

In the following, we discuss these issues by means of our grid models – all of them with shock emission described by our typical parameters ($T_s^{\infty} = 3 \cdot 10^6$ K and $f_X = 0.03$). In particular, we will provide estimates for suitable means of κ_{ν} , as a function of T_{eff} .

Figure 18 displays contour-plots of the radial dependence of the mass absorption coefficient in a D30 (upper panel) and in an S40 model (lower panel), as a function of wavelength. In accordance with our expectation from above, in both panels we note that κ_{ν} becomes constant when $r \gtrsim 1.2 R_*^{38}$ and the wavelength is lower than 20 Å (log $\lambda \leq 1.3$).

Longward of the O IV K-shell edge ($\lambda > 21$ Å), the radial variation of κ_{ν} depends on effective temperature and wind density. For the D30 model, κ_{ν} increases significantly with wavelength, but nevertheless does not vary with radius, because in this case the dominating ionization fraction of He II remains constant throughout the wind. In contrast, somewhat hotter models (e.g., D35), but particularly models with denser winds such as S40 display a different behavior. Here, the lower wind is dominated by He III, so that the background is weak, and one can already discriminate the C IV and C V edges around 10 R_* (indicated as dashed lines). Compared to the dwarf models, the total κ_{ν} in the inner wind is much lower, shows much more structure, and is influenced by the carbon and nitrogen opacities. Once helium begins to recombine in the outer wind, the background begins to dominate again, and the K-shell features vanish.

Fig. 19 illustrates the radial variation of the mass absorption coefficient for different wavelengths, and for our dwarf models with T_{eff} from 30 to 50 kK. Independent of T_{eff} , the radial variation of κ_{ν} is marginal at (and below) 10 Å. Around 20 Å, the variations in the inner/intermediate wind (until 10 R_*) are somewhat larger, due to changes in the oxygen ionization, where the specific positions of the corresponding edges play a role (see also Fig. 21). At 30 Å, we see a separation between D30 (black) with high values of κ_{ν} (He II dominating), hot models with low values of κ_{ν} (C v + low background, since helium completely ionized), and D35 (green) with a significantly varying κ_{ν} , due to the recombination of He III in the external wind. At 40 Å, finally, the behavior is similar, and only the κ_{ν} values for the cooler models are larger, because of the increasing He II background.

The analogous situation for supergiants is shown in Fig. 20. Whilst for dwarfs the variation of κ_v (when present) vanishes at around 10 R_* , here it is visible throughout the wind until large radii, for all but the coolest (black) and the hottest (red) model. Note that the limiting values (at the outermost radius) are similar to those of the corresponding dwarf models at $T_{\rm eff} = 30$ and 35 kK (recombined) and at $T_{\rm eff} = 50$ kK (He III). In contrast, for models with $T_{\rm eff} = 40$ and 45 kK the opacity continues to increase outwards, since the recombination is still incomplete.

Hervé et al. (2013) provided a similar figure to investigate the radial variation of κ_{ν} , in this case for a model of ζ Pup calculated by CMFGEN. Though the stellar parameters roughly agree with our S40 model, they considered a clumpy wind (with volume filling factor $f_V = 0.05$), and nuclear processed CNO abundances. Because this model shows an earlier recombination of helium, a larger nitrogen and a weaker oxygen K-shell edge, the actual values of κ_{ν} are somewhat different from our results (except at shortest wavelengths), but the basic trends are quite similar. In particular, our results support Hervé et al. (2013)'s idea of parameterizing the run of κ_{ν} : In any of the $\kappa_{\nu}(r)$ -curves displayed in Figs. 19 and 20, these curves either increase or slightly decrease, but eventually reach a plateau from a certain radius on (which differs for each model). This radius then separates two different regimes of κ_v that might be parameterized in an appropriate way (see Hervé et al. 2013 for details).

Instead of a parameterization, it is also possible to calculate meaningful averages of κ_{ν} and the corresponding scatter. The size of this scatter then allows us to conclude when (w.r.t. wavelength and $T_{\rm eff}$) a spatially constant mass absorption coefficient might be used to analyze X-ray line profiles. Instead of a straight average, we use here a density-weighted average (and a corresponding variance), to account for the fact that the optical depth, τ_{ν} , is the quantity which needs to be calculated with high precision:

$$\tau_{\nu} = \int_{R_{\min}}^{R_{\max}} \kappa_{\nu}(r)\rho(r)\mathrm{d}r =: \bar{\kappa}_{\nu} \int_{R_{\min}}^{R_{\max}} \rho(r)\mathrm{d}r \qquad \Rightarrow$$
$$\bar{\kappa}_{\nu} = \int_{R_{\min}}^{R_{\max}} \kappa_{\nu}(r)f(r)\mathrm{d}r, \qquad (11)$$

$$\operatorname{Var}(\kappa_{\nu}) = \int_{R_{\min}}^{R_{\max}} (\kappa_{\nu}(r) - \bar{\kappa}_{\nu})^2 f(r) \mathrm{d}r \qquad (12)$$

with p.d.f. $f(r)dr = \rho(r)dr / \left[\int_{R_{\min}}^{R_{\max}} \rho(r)dr\right]$.

In this approach³⁹, the density weights correspond to a probability distribution function (p.d.f.).

Fig. 21 displays such mean mass absorption coefficients, $\bar{\kappa}_{\nu}$, as a function of wavelength, averaged over the interval between 1.2 and 110.0 R_* , for our dwarf models.⁴⁰ The lower panel denotes the *relative* standard deviation, $\sqrt{\operatorname{Var}(\kappa_{\nu})}/\bar{\kappa}_{\nu}$. Also here, cold and hot models are clearly separated, with D35 in between (cf. with Fig. 19): for $\lambda \ge 21$ Å, the cold models are affected by a strong He II-background, whilst this background is weak for the hotter ones. In this 'long wavelength' region, the radial variation of κ_{ν} is large for model D35, due to recombining helium. There is also a considerable scatter between 18 and 21 Å, because of radial changes in the oxygen ionization. Overall, however, the assumption of a constant mass absorption coefficient (suitably averaged) is not too bad for the *complete* wavelength range (scatter below 20%), if we exclude model D35. Below 18 Å, the scatter becomes negligible, except at the Ne, Mg, and Si edges.

Even if $\kappa_v(r)$ can be approximated by a single number, $\bar{\kappa}_v$, the question is then about its value. For a comparison, the dashed line in Fig. 21 displays the (analytic) estimate, $\kappa_{\nu}^{\text{appr}}$ as provided by Eq. 9, using solar abundances and K-shell opacities only, with cross-sections from C IV, N IV, O IV, Ne IV, Mg IV, and Si IV. At least for the hotter models, this estimate is quite appropriate when comparing to the actual case, except for a somewhat erroneous description of the carbon edge(s): Since C v dominates in the hotter models and there is a ~ 4 Å difference between the C IV K-shell and the C v edge, this region is badly described by our approximation. For the cooler models, on the other hand, the difference between the dashed and the solid curves is (mostly) due to the helium background, which varies as a function of $T_{\rm eff}$, log g, and wind-density, thus affecting the actual value of $\bar{\kappa}_{\nu}$. Even below 18 Å, this background is still non-negligible for model D30, with a maximum deviation of roughly 30% close to the oxygen edge. Nevertheless, we conclude that for all dwarf

 $^{^{38}}$ to be on the safe side. In most cases, this limit – arising from fluctuations in the opacity background – is even lower.

³⁹ Note that the quantity R_{\min} indicates the lower boundary for the averaging process, and must not be confused with the onset radius of the X-ray emission.

⁴⁰ The impact of the chosen interval will be discussed below.

Fig. 19. Radial variation of the mass absorption coefficient in dwarf models, for specific values of wavelength. Black: $T_{\text{eff}} = 30$ kK, green: 35 kK; blue: 40 kK; magenta: 45 kK; red: 50 kK. All models calculated with $T_s^{\infty} = 3 \cdot 10^6$ K and $f_X = 0.03$. Note the different scales for κ_{ν} .

Fig. 20. As Fig. 19, but for supergiant models.

models with $T_{\rm eff} \ge 35$ kK the assumption of a constant mass absorption coefficient approximated by $\kappa_v^{\rm appr}$ is justified when $\lambda \le 18$ Å (at least within our present assumptions, i.e., solar abundances and unclumped winds with not too large optical depths, such that the averaging down to 1.2 R_* is reasonable; for different models, see below). In all other cases, results from NLTE-atmosphere modeling should be preferred.

The situation for our supergiant models is displayed in Fig. 22. Below 20 Å, the situation is similar to the dwarf case, though here the background is lower, even for the coolest model, and the approximation of $\bar{\kappa}_{\nu}$ by κ_{ν}^{appr} might now be applied at all temperatures. For $\lambda > 30$ Å, however, almost all models (except for S50) can no longer be described by a radially constant κ , since all models with $T_{\text{eff}} \leq 45$ kK show recombining helium of different extent, leading to strong variations throughout the wind.

Thus far, we considered models with solar abundances and unclumped winds. To illustrate the variation of the total and K-shell opacities with abundance (already investigated for particular models by, e.g., Cohen et al. 2010, 2014b), the dotted

Fig. 21. Upper panel: Density-weighted mean (Eq. 11) of the mass absorption coefficient, \bar{k}_{ν} , for the interval between 1.2 and 110 R_* , as a function of wavelengths, and for dwarf models with $T_s^{\infty} = 3 \cdot 10^6$ K and $f_X = 0.03$. Solar abundances following Asplund et al. (2009) have been adopted. Dashed: approximate, radius-independent κ_{ν}^{appr} (Eq. 9), using solar abundances and K-shell opacities only, with cross-sections from C rv (with threshold at 35.7 Å), N rv (27.0 Å), O rv (20.8 Å), Ne rv (13.2 Å), Mg rv (9.0 Å), and Si rv (6.4 Å). The C v edge (at 31.6 Å) appears as unresolved in our frequency grid. Dotted: as dashed, but with nuclear processed CNO abundances as derived for ζ Pup by Bouret et al. (2012). Note that the nitrogen abundance is more than a factor of 10 larger than the solar one. Dashed and dotted lines serve also as a guideline for comparison with similar figures.

Lower panel: Relative standard deviation, $\sqrt{Var(\kappa_v)}/\bar{\kappa}_v$ (see Eq. 12), for the same models. The dotted line denotes a relative scatter of 15%.

Fig. 22. As Fig. 21, but for supergiant models.

lines in Figs. 21 and 22 denote the approximate K-shell opacities, κ_v^{appr} , for the case of highly processed CNO material, based on the abundances derived for ζ Pup by Bouret et al. (2012). Here, the carbon and oxygen abundances are depleted by 0.8 and 0.6 dex, respectively, whilst the nitrogen abundance is extremely enhanced (by ~ 1.3 dex), compared to the solar values. Such a composition leads to weak C and O K-shell edges, but to an enormous nitrogen edge (dotted vs. dashed line).

Now, if the individual abundances are known during an analysis, there will be no problem, and κ_{ν} might be approximated by either $\kappa_{\nu}^{\text{appr}}$ below 18 Å or calculated by means of NLTEmodel atmospheres, simply accounting for these abundances. However, considerable uncertainties even in the low wavelength regime might result when the abundances are not known. From comparing the dashed and the dotted line, we estimate this uncertainty as roughly 50% for $\bar{\kappa}_{\nu}$, and thus for τ_{ν} and \dot{M} (when the mass-loss rate shall be derived). A similar value has already been estimated by Cohen et al. (2014b). In the range between the oxygen and the carbon edge (20 to 35 Å), the situation is even worse, and we conclude that the corresponding absorption coefficients are prone to extreme uncertainties when the abundances have to be adopted without further verification. In particular, getting κ_{ν} right around 25 Å is important for measuring the N emission lines at and near that wavelength (e.g., N vi 24.9, N vII 24.78), and thus measuring the N abundance directly. But at longer wavelengths, where κ_{ν} will vary even more strongly with radius, and even though nitrogen emission lines are not directly affected, the (direct) ionization of CNO etc. is affected, and so optical and UV line strengths are affected too, as discussed in the previous sections.

The impact of clumping is less severe. Comparing Fig. C.1 (Appendix) with Figs. 21 and 22, we see that models accounting for optically thin clumping ('micro-clumping') with typical clumping factors⁴¹ and adequately reduced mass-loss rates give rather similar results compared with unclumped models. Again, the scatter of κ_{ν} is negligible below 18 Å. 'Only' the region longward of 20 Å is stronger contaminated by the He II background, since the clumped models recombine earlier than the unclumped ones. Note that the K-shell mass absorption coefficients themselves are *not* affected by optically thin clumping, since the opacities scale linearly with density.

Finally, Fig. C.2 (Appendix) investigates the consequences of averaging κ_{ν} in the outer wind alone (in the interval between 10 and 110 R_*), which would be adequate if the wind would become optically thick at such radii (which for short wavelengths and O-star winds is quite unlikely, because of the low value of κ_{ν}). Anyway, below 18 Å the differences to the original values are small. Note that the hot dwarf models now behave almost exactly as estimated by $\kappa_{\nu}^{\text{appr}}$, because He II vanishes in the outer regions of these objects.

Further conclusions on this topic are provided in the next section.

6. Summary and conclusions

In this paper, we described the implementation of X-ray emission from wind-embedded shocks into the unified, NLTE atmosphere/spectrum synthesis code FASTWIND, discussed various tests, and presented some first result.

Our implementation follows closely corresponding work by Pauldrach et al. (2001) (for WM-BASIC), which in turn is based on the shock cooling zone model developed by Feldmeier et al. (1997a), with the additional possibility to consider isothermal shocks. The (present) description of the shock-distribution and strength is provided by four input, 'X-ray emission parameters', controlling the filling factor, the run of the shock temperature, and the radial onset of the emitting plasma. We account for Kshell absorption and Auger ionization, allowing for more than one final ionization stage due to cascade ionization processes. Most of our test calculations are based on a grid of 11 models (supergiants and dwarfs within T_{eff} = 30 to 55 kK), each of them with 9 different X-ray emission parameter sets, but many more models have been calculated for various comparisons, including models with optically thin clumping.

A first test investigated the reaction when varying important X-ray emission parameters. For radially increasing shockstrengths, the emergent flux remains almost unaffected if the onset radius is lowered compared to its default value (roughly 1.5 R_*), whilst increasing the onset has a considerable effect in the range between \sim 350 Å and at least the He II edge. Filling factor and maximum shock temperature affect the ionization fractions, particularly of the highly ionized species. We confirm some earlier predictions for scaling relations for X-ray luminosities (as a function of M/v_{∞}) in the case of optically thin and thick continua⁴², but we noted that for our hottest models these luminosities can become contaminated by 'normal' stellar radiation, for energies below ~150 eV. Thus, we suggested to choose a lower integration limit of 0.15 keV (or even 0.3 keV, to be on the safe side) when comparing the X-ray luminosities of different stars or theoretical models. Finally, we found an excellent agreement between FASTWIND and WM-BASIC fluxes, demonstrating a similar ionization balance, and a satisfactory agreement between corresponding X-ray luminosities. Overall, the impact of typical shock emission affects the radiation field in the wind for all wavelengths $\lambda < 350$ Å, thus modifying all photo-rates for ions with ionization edges in this regime.

Investigating the ionization fractions within our model grid allowed us to study the impact of shock radiation for the proper description of important ions, i.e., those with meaningful wind lines (e.g., C IV, N IV, N V, O V, O VI, Si IV, and P V). If we denote models with T_{eff} = 30 to 35 kK as 'cool', models with T_{eff} = 35 to 45 kK as 'intermediate' and models with T_{eff} = 45 to 55 kK as 'hot' (note the overlap), we can summarize our findings as follows. *Not* (or only marginally) affected by shock emission (with typical parameters and our parameterization of the shockstrengths) are

- in dwarfs: C III, C IV, N III (cool), N IV (cool), O IV (intermediate), Si IV, P V (cool+intermediate)
- in supergiants: C III (hot), C IV (hot), N IV (cool), O IV (intermediate), Si IV (hot).

In almost all other cases, lower stages (C III, C IV, N III, N IV, O IV (hot), Si IV, and P V) are depleted, i.e., corresponding wind lines become weaker, and higher stages (N v, O IV (cool), O v, O VI) become enhanced, i.e., corresponding wind lines become stronger when accounting for shock emission.

We studied in some detail how the ionization fractions change when the two most important parameters, filling factor and maximum shock temperature, are varied. For most ions, the filling factor has a larger influence than T_s^{∞} , but particularly O vi and P v (the latter only for higher filling factors and shock temperatures) show a strong reaction to both parameters.

Due to the importance of P v with respect to mass-loss and wind-structure diagnostics, we re-investigated its behavior, and confirm previous results that for typical X-ray emission parameters this ion is only weakly or moderately affected (by factors of two for intermediate and hot supergiants at $v(r)/v_{\infty} = 0.5$ and by factors of 10 at $v(r)/v_{\infty} = 0.8$), though for a strong X-ray radiation field the depletion can reach much higher factors. A com-

⁴¹ $f_{\rm cl} = 20$ corresponding to a volume filling factor, $f_{\rm V} = 0.05$

⁴² though a discrepancy with recent work by Owocki et al. (2013) was identified, which needs to be investigated further

parison of P v ionization fractions with results from CMFGEN (Bouret et al. 2012) provided a reasonable agreement.

Not only metals, but also He can be affected by shock emission, due to the location of the He II edge and He II 303 in the EUV. Significant effects, however, have only been found in the winds of cool supergiants, where particularly He II 1640 (emission and high-velocity absorption) and He II 4686 (emission) become stronger, due to increased recombination cascades and increased pumping of the n = 2 level in case of He II 1640.

When comparing our ionization fractions with those calculated by WM-BASIC, we found a good, though not perfect, agreement, which turned out to be true also for various UV-lines. When comparing with Krtička & Kubát (2009), on the other hand, a similar agreement over the complete covered temperature range was found only for few ions; for the majority, such agreement is present only at specific temperatures.

It is well known that Auger ionization can play an important role for the ionization balance of specific ions. To further investigate this issue, we compared the ionization fractions of all ions considered in this study when including (default) or excluding this process in our NLTE treatment. Overall, it turned out that only N v1 and O v1 (as previously known) are significantly affected by Auger ionization, but, at least in our models (with radially increasing shock temperatures), only in the outer wind. For the inner and intermediate wind, direct EUV/XUV ionization due to shock emission dominates, which is generally true for all other considered ions.⁴³

As an interesting by-product of our investigation, it turned out that dielectronic recombination of O v can have a considerable influence on the ionization balance of oxygen (O v vs. O v), particularly for dwarfs around 45 kK.

In the last part of this paper, we provided an extensive discussion of the (high-energy) mass absorption coefficient, κ_{γ} , regarding its spatial variation and dependence on T_{eff} . This topic is particularly relevant for various approaches to analyze X-ray emission lines. To summarize and conclude, we found that (i) the approximation of a radially constant κ_{γ} can be justified for $r \ge 1.2R_*$ and $\lambda \le 18$ Å, and also for many models at longer wavelengths. (ii) In order to estimate the actual value of this quantity, however, the He II background⁴⁴ needs to be considered from detailed modeling, at least for wavelengths longer than 18 to 20 Å. Moreover, highly processed CNO material can change the actual value of κ_{γ} considerably, particularly for $\lambda \ge 20$ Å, and estimates for the optical depth, τ_{γ} , become highly uncertain in this regime if the individual abundances are unknown.

In this context, it is reassuring to note that, e.g., the mass-loss determinations by Cohen et al. (2014b) using X-ray line spectroscopy (via determining the optical depths of the cool wind material, under the assumption of spatially constant κ_v) rely on 16 lines observed by CHANDRA, where 14 out of these 16 lines are shortward of 19 Å. The issues summarized above will be a much bigger problem for O vII and nitrogen X-ray emission line measurements (O vII at 21.6-22.1 Å, N vII at 24.78 Å and N vI at 24.9 Å) that are planned to independently constrain, with high precision, the nitrogen/oxygen content in (a few) massive O-stars (Leutenegger et al. 2013a). To this end, a detailed modeling of κ_v (particularly regarding the helium ionization) will certainly be advisable for such an analysis.

Having finalized and carefully tested our implementation of emission from wind-embedded shocks, we are now in a position to continue our work on the quantitative spectroscopy of massive stars. As outlined in the introduction, we will concentrate on determining the carbon and oxygen abundances in O- and early B-stars observed during the two VLT-FLAMES surveys conducted within our collaboration, by means of optical and, when available, UV spectroscopy. During such an analysis, the X-ray emission parameters need to be derived in parallel with the other, main diagnostics, at least in principle. We then have to check in how far the derived abundances depend on corresponding uncertainties.

Note further that any such UV analysis also needs to consider the effects of optically thick clumping (e.g., Oskinova et al. 2007, Sundqvist et al. 2011, Šurlan et al. 2013, Sundqvist et al. 2014). In parallel with the implementation of wind-embedded shocks presented here, we have updated FASTWIND to also account properly for such optically thick clumping (porosity in physical and velocity space), following Sundqvist et al. (2014); these models will be presented in an upcoming (fourth) paper of this series.

Regarding quantitative spectroscopic studies accounting for X-ray ionization effects, the parameterization represented by Eq. 7 is certainly not the final truth, and actually also not the best encapsulation of the results from present-day numerical simulations. Though this probably does not matter too much for most applications, it might be worth thinking about a better representation, and how our results would change if the stronger and weaker shocks were allowed to be more spatially mixed.

LDI simulations (e.g., Feldmeier et al. 1997b. Dessart & Owocki 2003, Sundqvist & Owocki 2013) indicate that the velocity dispersion peaks quite close to R_{\min} (~ 1.5-2.0 R_*) and then falls off. And the same simulations show also some strong shocks near R_{\min} . From the observational side, f/i ratios of ions that form at higher temperatures (e.g., Si xIII) indicate a substantial amount of high-temperature plasma (~10⁷ K) near R_{min} (e.g., Waldron & Cassinelli 2001, Waldron & Cassinelli 2007), and Leutenegger et al. (2006) found an onset radius of $1.1^{+0.4/-0.1} R_*$ for the S xv line. On the other hand, Cohen et al. (2014a) showed that the shock temperature distribution is very strongly skewed toward weak shocks, and our parameterization Eq. 7 allows us to include that feature already now.

Acknowledgements. The authors like to thank the referee, David Cohen, for helpful comments and suggestions. LPC gratefully acknowledges support from the Brazilian Coordination for the Improvement of Higher Education Personnel (CAPES), under grant 0964-13-1. JOS acknowledges funding from the European Union's Horizon 2020 research and innovation programme under the Marie Sklodowska-Curie grant agreement No 656725. Many thanks also to J.-C. Bouret for providing us with the ionization fractions of P tv and P v from 'his' models of HD 163758 and HD 16691.

References

Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. 2009, ARA&A, 47, 481 Bouret, J.-C., Hillier, D. J., Lanz, T., & Fullerton, A. W. 2012, A&A, 544, A67

- Bruccato, R. J. & Mihalas, D. 1971, MNRAS, 154, 491
- Butler, K. & Giddings, J. R. 1985, Newsl. Anal. Astron. Spectra, 9
- Cassinelli, J. & Olson, G. 1979, ApJ, 229, 304
- Cassinelli, J. P., Cohen, D. H., Macfarlane, J. J., et al. 1995, ApJ, 438, 932
- Cassinelli, J. P., Cohen, D. H., Macfarlane, J. J., Sanders, W. T., & Welsh, B. Y. 1994, ApJ, 421, 705
- Cassinelli, J. P. & Swank, J. H. 1983, ApJ, 271, 681
- Chlebowski, T., Harnden, F., & Sciortino, S. 1989, ApJ, 341, 427
- Cohen, D. H., Cassinelli, J. P., & MacFarlane, J. J. 1997, ApJ, 487, 867
- Cohen, D. H., Cooper, R. G., Macfarlane, J. J., et al. 1996, ApJ, 460, 506
- Cohen, D. H., Gagné, M., Leutenegger, M. A., et al. 2011, MNRAS, 415, 3354

 $^{^{43}}$ In clumped winds, the presence of a low-density interclump medium is essential for the formation of O vi (Zsargó et al. 2008).

 $^{^{\}rm 44}\,$ and, to a lesser extent, also the bound-free background from highly abundant metals

- Cohen, D. H., Kuhn, M. A., Gagné, M., Jensen, E. L. N., & Miller, N. A. 2008, MNRAS, 386, 1855
- Cohen, D. H., Leutenegger, M. A., Wollman, E. E., et al. 2010, MNRAS, 405, 2391
- Cohen, D. H., Li, Z., Gayley, K. G., et al. 2014a, MNRAS, 444, 3729
- Cohen, D. H., Wollman, E. E., Leutenegger, M. A., et al. 2014b, MNRAS, 439, 908
- Crowther, P. A., Hillier, D. J., Evans, C. J., et al. 2002, ApJ, 579, 774
- Daltabuit, E. & Cox, D. 1972, ApJ, 177, 855
- Dessart, L. & Owocki, S. P. 2003, A&A, 406, L1
- Evans, C., Hunter, I., Smartt, S., et al. 2008, The Messenger, 131, 25
- Evans, C. J., Taylor, W. D., Hénault-Brunet, V., et al. 2011, A&A, 530, A108 Feldmeier, A. 1995, A&A, 299, 523
- Feldmeier, A., Kudritzki, R.-P., Palsa, R., Pauldrach, A. W. A., & Puls, J. 1997a, A&A, 320, 899
- Feldmeier, A., Puls, J., & Pauldrach, A. W. A. 1997b, A&A, 322, 878
- Fullerton, A. W., Massa, D. L., & Prinja, R. K. 2006, ApJ, 637, 1025
- Gabler, R., Gabler, A., Kudritzki, R. P., Puls, J., & Pauldrach, A. 1989, A&A, 226, 162
- Garcia, M. 2005, PhD thesis, University of La Laguna (Teneriffe)
- Giddings, J. R. 1981, PhD thesis, , University of London, (1981)
- Gräfener, G., Koesterke, L., & Hamann, W.-R. 2002, A&A, 387, 244
- Groenewegen, M. A. T. & Lamers, H. J. G. L. M. 1989, A&AS, 79, 359
- Hamann, W.-R. & Oskinova, L. 2012, in COSPAR Meeting, Vol. 39, 39th COSPAR Scientific Assembly, 716
- Harnden, Jr., F. R., Branduardi, G., Gorenstein, P., et al. 1979, ApJ, 234, L51
- Haser, S. M. 1995, PhD thesis, Ludwig-Maximilians-Universität München
- Hauschildt, P. H. 1992, Journal of Quantitative Spectroscopy and Radiative Transfer, 47, 433
- Hervé, A., Rauw, G., & Nazé, Y. 2013, A&A, 551, A83
- Hillier, D. J., Kudritzki, R. P., Pauldrach, A. W., et al. 1993, A&A, 276, 117
- Hillier, D. J. & Miller, D. L. 1998, ApJ, 496, 407
- Hubeny, I. 1998, in Astronomical Society of the Pacific Conference Series, Vol. 138, 1997 Pacific Rim Conference on Stellar Astrophysics, ed. K. L. Chan, K. S. Cheng, & H. P. Singh, 139
- Huenemoerder, D. P., Oskinova, L. M., Ignace, R., et al. 2012, ApJ, 756, L34
- Hunter, I., Brott, I., Lennon, D. J., et al. 2008, ApJ, 676, L29
- Hunter, I., Dufton, P. L., Smartt, S. J., et al. 2007, A&A, 466, 277
- Kaastra, J. & Mewe, R. 1993, A&A, 97, 443
- Krolik, J. & Raymond, J. 1985, ApJ, 298, 660
- Krtička, J. & Kubát, J. 2001, A&A, 369, 222
- Krtička, J. & Kubát, J. 2009, MNRAS, 394, 2065
- Krtička, J. & Kubát, J. 2012, MNRAS, 427, 84
- Kubát, J., Puls, J., & Pauldrach, A. W. A. 1999, A&A, 341, 587
- Kudritzki, R.-P. & Puls, J. 2000, ARA&A, 38, 613
- Lamers, H. & Rogerson, J. 1978, A&A, 66, 417
- Lamers, H. J. G. L. M., Cerruti-Sola, M., & Perinotto, M. 1987, ApJ, 314, 726
- Lamers, H. J. G. L. M. & Morton, D. C. 1976, ApJS, 32, 715
- Leutenegger, M. A., Cohen, D., Neely, J., Owocki, S., & Sundqvist, J. 2013a, in Massive Stars: From alpha to Omega, 42
- Leutenegger, M. A., Cohen, D. H., Sundqvist, J. O., & Owocki, S. P. 2013b, ApJ, 770.80
- Leutenegger, M. A., Paerels, F. B. S., Kahn, S. M., & Cohen, D. H. 2006, ApJ, 650, 1096
- Lucy, L. B. 1982, ApJ, 255, 278
- Lucy, L. B. & Solomon, P. M. 1970, ApJ, 159, 879
- Macfarlane, J. J., Cohen, D. H., & Wang, P. 1994, A&A, 437, 351
- Macfarlane, J. J., Waldron, W. L., Corcoran, M. F., et al. 1993, ApJ, 419, 813
- Martins, F., Escolano, C., Wade, G. A., et al. 2012, A&A, 538, A29
- Martins, F., Hervé, A., Bouret, J.-C., et al. 2015a, A&A, 575, A34
- Martins, F. & Hillier, D. J. 2012, A&A, 545, A95
- Martins, F., Simón-Díaz, S., Palacios, A., et al. 2015b, A&A, 578, A109
- Mihalas, D. & Hummer, D. G. 1973, ApJ, 179, 827
- Nazé, Y., Broos, P. S., Oskinova, L., et al. 2011, ApJS, 194, 7
- Nazé, Y., Flores, C. A., & Rauw, G. 2012, A&A, 538, A22
- Nazé, Y., Oskinova, L. M., & Gosset, E. 2013, ApJ, 763, 143
- Olson, G. & Castor, J. 1981, ApJ, 244, 179
- Oskinova, L. M., Feldmeier, A., & Hamann, W.-R. 2006, MNRAS, 372, 313
- Oskinova, L. M., Hamann, W.-R., & Feldmeier, A. 2007, A&A, 476, 1331
- Owocki, S. P. 1994, Ap&SS, 221, 3
- Owocki, S. P., Castor, J. I., & Rybicki, G. B. 1988, ApJ, 335, 914
- Owocki, S. P. & Cohen, D. H. 1999, ApJ, 520, 833
- Owocki, S. P. & Cohen, D. H. 2006, ApJ, 648, 565
- Owocki, S. P. & Rybicki, G. B. 1984, ApJ, 284, 337
- Owocki, S. P., Sundqvist, J. O., Cohen, D. H., & Gayley, K. G. 2013, MNRAS, 429.3379
- Pauldrach, A. W. A., Hoffmann, T. L., & Lennon, M. 2001, A&A, 375, 161
- Pauldrach, A. W. A., Kudritzki, R. P., Puls, J., & Butler, K. 1990, A&A, 228,

- 125
- Pauldrach, A. W. A., Kudritzki, R. P., Puls, J., Butler, K., & Hunsinger, J. 1994, A&A, 283, 525
- Przybilla, N., Firnstein, M., Nieva, M. F., Meynet, G., & Maeder, A. 2010, A&A, 517 A38
- Puls, J. 2009, Communications in Asteroseismology, 158, 113
- Puls, J., Kudritzki, R.-P., Herrero, A., et al. 1996, A&A, 305, 171
- Puls, J., Owocki, S. P., & Fullerton, A. W. 1993, A&A, 279, 457
- Puls, J., Urbaneja, M. A., Venero, R., et al. 2005, A&A, 435, 669
- Rauw, G., Hervé, A., Nazé, Y., et al. 2015, A&A, 580, A59
- Raymond, J. & Smith, B. 1977, ApJS, 35, 419
- Rivero González, J. G., Puls, J., & Najarro, F. 2011, A&A, 536, A58
- Rivero González, J. G., Puls, J., Najarro, F., & Brott, I. 2012a, A&A, 537, A79
- Rivero González, J. G., Puls, J., Najarro, F., & Massey, P. 2012b, A&A, 543, A95
- Sana, H., Rauw, G., Naze, Y., Gosset, E., & Vreux, J.-M. 2006, MNRAS, 372, 661
- Seaton, M. J. 1958, MNRAS, 118, 504
- Seward, F. D., Forman, W. R., Giacconi, R., et al. 1979, ApJ, 234, L55
- Simon, M. & Axford, W. 1966, Planet.Space Sci., 14, 901
- Smith, R. K., Brickhouse, N. S., Liedahl, D. A., & Raymond, J. C. 2001, ApJ, 556, L91
- Snow, T. & Morton, D. 1976, ApJS, 32, 429
- Sundqvist, J. O. & Owocki, S. P. 2013, MNRAS, 428, 1837
- Sundqvist, J. O., Owocki, S. P., Cohen, D. H., Leutenegger, M. A., & Townsend, R. H. D. 2012a, MNRAS, 420, 1553
- Sundqvist, J. O., Owocki, S. P., & Puls, J. 2012b, in Astronomical Society of the Pacific Conference Series, Vol. 465, Proceedings of a Scientific Meeting in Honor of Anthony F. J. Moffat, ed. L. Drissen, C. Robert, N. St-Louis, & A. F. J. Moffat. 119
- Sundqvist, J. O., Puls, J., Feldmeier, A., & Owocki, S. P. 2011, A&A, 528, A64
- Sundqvist, J. O., Puls, J., & Owocki, S. P. 2014, A&A, 568, A59
- Trundle, C., Lennon, D. J., Puls, J., & Dufton, P. L. 2004, A&A, 417, 217
- Šurlan, B., Hamann, W.-R., Aret, A., et al. 2013, A&A, 559, A130
- Verner, D. A. & Yakovlev, D. G. 1995, A&AS, 109, 125
- Walborn, N. R. & Panek, R. J. 1984, ApJ, 280, L27
- Waldron, W. L. & Cassinelli, J. P. 2007, ApJ, 668, 456
- Waldron, W. L. & Cassinelli, J. P. 2010, A&A, 711, L30
- Waldron, W. L. & Cassinelli, J. P. 2001, ApJ, 548, L45
- Zhekov, S. A. & Palla, F. 2007, MNRAS, 382, 1124
- Zsargó, J., Hillier, D. J., Bouret, J.-C., et al. 2008, ApJ, 685, L149

Appendix A: Ionization fractions of selected ions: Dependence on X-ray filling factor and shock temperature

Figures A.1 to A.10 display the reaction of C IV, N V, O V, O VI, and P v on varying the X-ray filling factors and shock temperatures within our supergiant and dwarf models, as a function of $T_{\rm eff}$. For further explanation and discussion, see Sect. 5.1.3.

Appendix B: Comparison with WM-BASIC: Ionization fractions and UV line-profiles

In Figs. B.1 and B.2 we compare the ionization fractions of specific ions, as calculated by FASTWIND and WM-BASIC, for dwarf and supergiant models, respectively. Fig. B.3 compares corresponding strategic UV-line profiles for NIV 1720, N v 1238,1242, O v 1371, O VI 1031,1037, and P v 1117,1128. Further explanation and discussion is provided in Sect. 5.1.4.

Fig. A.1. Ionization fractions of C IV (at $v(r) = 0.5v_{\infty}$), as a function of T_{eff} , and for different X-ray emission parameters. Solid: supergiant models; dashed: dwarf models; Black: models with shock emission; magenta: models without shock emission. *For clarity, the ionization fractions of dwarf models have been shifted by one dex.*

Fig. A.2. Left panel - As above (C iv at $v(r) = 0.5v_{\infty}$), but now for dwarf models alone and for all X-ray emission parameters included in our grid. Note that the fractions have *not* been shifted here. *Right panel* - as left, but for supergiant models.

Fig. A.3. As Fig. A.1, but for N v at $v(r) = 0.6v_{\infty}$.

Fig. A.4. As Fig. A.2, but for N v ($v(r) = 0.6v_{\infty}$).

Fig. A.5. As Fig. A.1, but for O v at $v(r) = 0.6v_{\infty}$.

Fig. A.6. As Fig. A.2, but for O v ($v(r) = 0.6v_{\infty}$).

Fig. A.7. As Fig. A.1, but for O vI at $v(r) = 0.6v_{\infty}$.

Fig. A.8. As Fig. A.2, but for O v ($v(r) = 0.6v_{\infty}$).

Fig. A.9. As Fig. A.1, but for P v at $v(r) = 0.5v_{\infty}$.

Fig. A.10. As Fig. A.2, but for $Pv(v(r) = 0.5v_{\infty})$.

Fig. B.1. Ionization fractions of specific ions, as calculated by FASTWIND (black) and WM-BASIC (magenta), for our dwarf models and as a function of T_{eff} . If not stated explicitly inside the individual panels, the fractions have been evaluated at $v(r) = 0.5v_{\infty}$. See Sect. 5.1.4.

Fig. B.2. As Fig. B.1, but for supergiant models.

Fig. B.3. Emergent line profiles for strategic UV lines (N tv 1720, N v 1238,1242, O v 1371, O v1 1031,1037, and P v 1117,1128), as calculated by WM-BASIC (green) and FASTWIND (black), for models S30 (top), D40, S40, D50, and S50 (bottom). All profiles have been calculated with a radially increasing microturbulence, with maximum value $v_{turb}(max) = 0.1v_{\infty}$, and have been convolved with a typical rotation velocity, $v \sin i = 100 \text{ km s}^{-1}$. The absorption feature between the two P v components is due to Si tv 1122. See Sect. 5.1.4.

Appendix C: Averaged mass absorption coefficients - clumped winds and dependence on averaging interval

Fig. C.1 displays the density-weighted mean (Eq. 11) of the mass absorption coefficient as a function of wavelength, for dwarf (left) and supergiant (right) models. The figure has a similar layout as Figs. 21 and 22, but has been calculated for clumped models ($f_{cl} = 20$), and mass-loss rates reduced by a factor of $\sqrt{20}$. Fig. C.2 is also analogous to Figs. 21 and 22, but now the absorption coefficient has been averaged over the interval between 10 and 110 R_* . For details and discussion, see Sect. 5.4.

Fig. C.1. As Figs. 21 and 22, but for clumped models with $f_{cl} = 20$ (corresponding to $f_V = 0.05$) and mass-loss rates reduced by a factor of $\sqrt{20}$. Left: dwarf models; right: supergiant models.

Fig. C.2. As Figs. 21 and 22, but averaged over the interval between 10 and 110 R_{*}. Left: dwarf models; right: supergiant models.